

Date of Deposit: 9/11/2000

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of: Hong Zhang and Andrew T. Watt

For: Antisense Modulation of caspase 7 Expression

BOX SEQUENCE

Assistant Commissioner for Patents

Washington DC 20231

## PATENT APPLICATION TRANSMITTAL LETTER

Transmitted herewith for filing, please find the following:

- The specification of the above-referenced patent application;
- An **executed** Declaration or Oath and Power of Attorney;
- An Assignment of the invention to **Isis Pharmaceuticals Inc.** with recordation cover sheet (PTO Form PTO-1595) and \$40.00 cover fee;
- An **executed** Verified Statement Claiming Small Entity Status under 37 CFR 1.9 and 1.27;
- Statement to Support Filing and Submission of DNA/Amino Acid Sequences in Accordance with 37 CFR § 1.821 through 1.825;
- Sequence listing in computer readable form in accordance with 37 C.F.R. § 1.821(e); and
- An Information Disclosure Statement with references.

The filing fee has been calculated as shown below:

For:	No. Filed	No. Extra	Rate	Fee
BASE FEE				\$345.00
Total Claims	20 - 20 =	0	X \$9=	\$0
Indep. Claims	2 - 3 =	0	x\$39=	\$0
TOTAL				\$ 345.00

The Commissioner is hereby authorized to charge the following fees to Deposit Account No. 500252:

- the amount of \$385.00 for the above listed fees;
- payment of the following fees associated with this communication or credit any overpayment;
- any additional filing fees required under 37 CFR 1.16 including fees for presentation of extra claims; and
- any additional patent application processing fees under 37 CFR 1.17 and under 37 CFR 1.20 (d).

Triplicate copies of this transmittal are enclosed.

Date: Sept. 11 2000

*Laurel Spear Bernstein*  
 Laurel Spear Bernstein  
 Registration No. 37,280  
 Isis Pharmaceuticals, Inc.

Please address all correspondence to:

Jane Massey Licata  
 Law Offices of Jane Massey Licata  
 66 East Main Street  
 Marlton NJ 08053  
 (856) 810-1515

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of: Hong Zhang and Andrew T. Watt

For: Antisense Modulation of caspase 7 Expression

BOX SEQUENCE

Assistant Commissioner for Patents  
Washington DC 20231PATENT APPLICATION TRANSMITTAL LETTER

Transmitted herewith for filing, please find the following:

- The specification of the above-referenced patent application;
- An executed Declaration or Oath and Power of Attorney;
- An Assignment of the invention to Isis Pharmaceuticals Inc. with recordation cover sheet (PTO-Form PTO-1595) and \$40.00 cover fee;
- An executed Verified Statement Claiming Small Entity Status under 37 CFR 1.9 and 1.27;
- Statement to Support Filing and Submission of DNA/Amino Acid Sequences in Accordance with 37 CFR \$ 1.821 through 1.825;
- Sequence listing in computer readable form in accordance with 37 C.F.R. \$ 1.821(e); and
- An Information Disclosure Statement with references.

The filing fee has been calculated as shown below:


For:	No. Filed	No. Extra	Rate	Fee
BASE FEE				\$345.00
Total Claims	20 - 20 =	0	X \$9=	\$0
Indep. Claims	2 - 3 =	0	x\$39=	\$0
TOTAL				\$ 345.00

The Commissioner is hereby authorized to charge the following fees to Deposit Account No. 500252:

- the amount of \$385.00 for the above listed fees;
- payment of the following fees associated with this communication or credit any overpayment;
- any additional filing fees required under 37 CFR 1.16 including fees for presentation of extra claims; and
- any additional patent, application processing fees under 37 CFR 1.17 and under 37 CFR 1.20 (d).

Triplicate copies of this transmittal are enclosed.

Date: Sept. 11 2000

  
Laurel Spear Bernstein  
Registration No. 37,280  
Isis Pharmaceuticals, Inc.

Please address all correspondence to:

Jane Massey Licata  
Law Offices of Jane Massey Licata  
66 East Main Street  
Marlton NJ 08053  
(856) 810-1515

Date of Deposit: 9/11/2000

Applicant or Patentee: Hong Zhang and Andrew T. Watt

Serial or Patent No.: not yet assigned

Date Filed or Issued: herewith

For: Antisense Modulation of caspase 7 Expression

**VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS  
(37 CFR 1.9(f) and 1.27(c) - SMALL BUSINESS CONCERN)**

I hereby declare that I am an official empowered to act on behalf of the concern identified below:

**NAME OF CONCERN:** Isis Pharmaceuticals Inc.  
**ADDRESS OF CONCERN:** 2292 Faraday Ave  
 Carlsbad, California 92008

I hereby declare that the above-identified small business concern qualifies as a small business concern as defined in 13 CFR 121.12, and reproduced in 37 CFR 1.9(d), for purposes of paying reduced fees under section 41(a) and (b) of Title 35, United States Code, in that: (1) the number of employees of the concern, including those of its affiliates, does not exceed 500 persons; and (2) the concern has not assigned, granted, conveyed, or licensed, and is under no obligation under contract or law to assign, grant, convey, or license, any rights in the invention to any person who could not be classified as an independent inventor if that person had made the invention, or to any concern which would not qualify as a small business concern or a nonprofit organization under this section.

For purposes of this statement, (1) the number of employees of the business concern is the average over the previous fiscal year of the concern of the persons employed on a full-time, part-time or temporary basis during each of the pay periods of the fiscal year, and (2) concerns are affiliates of each other when either, directly or indirectly, one concern controls or has the power to control the other, or a third party or parties controls or has the power to control both.

I hereby declare that rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the invention entitled **Antisense Modulation of caspase 7 Expression** by inventor(s) Hong Zhang and Andrew T. Watt described in the specification filed herewith.

If the rights held by the above-identified small business concern are not exclusive, each individual, concern or organization having rights in the invention is listed below\* and no rights to the invention are held by any person, other than the inventor, who would not qualify as an independent inventor under 37 CFR 1.9(c) if that person made the invention, or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d), or a nonprofit organization under 37 CFR 1.9(e).

\*NOTE: Separate verified statements are required for each named person, concern or organization having rights to the invention averring to their status as small entities.  
 (37 CFR 1.27)

**FULL NAME:** ( ) INDIVIDUAL ( ) SMALL BUSINESS CONCERN  
 ( ) NONPROFIT ORGANIZATION

**ADDRESS:**

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b))

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

**NAME OF PERSON SIGNING**  
**TITLE OF PERSON SIGNING**  
**ADDRESS OF PERSON SIGNING**

C. Frank Bennett  
 Vice President, Biology  
 2292 Faraday Avenue  
 Carlsbad, CA 92008

*C. Frank Bennett*  
 SIGNATURE

*9-6-00*  
 DATE

5                   **ANTISENSE MODULATION OF CASPASE 7 EXPRESSION****FIELD OF THE INVENTION**

10           The present invention provides compositions and methods  
for modulating the expression of caspase 7. In particular,  
this invention relates to compounds, particularly  
oligonucleotides, specifically hybridizable with nucleic  
acids encoding caspase 7. Such compounds have been shown to  
15   modulate the expression of caspase 7.

**BACKGROUND OF THE INVENTION**

          Apoptosis, or programmed cell death, is a naturally  
20   occurring process that has been strongly conserved during  
evolution to prevent uncontrolled cell proliferation. This  
form of cell suicide plays a crucial role in ensuring the  
development and maintenance of multicellular organisms by  
eliminating superfluous or unwanted cells. However, if this  
25   process goes awry becoming overstimulated, cell loss and  
degenerative disorders including neurological disorders such  
as Alzheimers, Parkinsons, ALS, retinitis pigmentosa and  
blood cell disorders can result. Stimuli which can trigger  
apoptosis include growth factors such as tumor necrosis  
30   factor (TNF), Fas and transforming growth factor beta (TGF $\beta$ ),  
neurotransmitters, growth factor withdrawal, loss of  
extracellular matrix attachment and extreme fluctuations in  
intracellular calcium levels (Afford and Randhawa, *Mol.*  
*Pathol.*, 2000, 53, 55-63).

Alternatively, insufficient apoptosis, triggered by growth factors, extracellular matrix changes, CD40 ligand, viral gene products neutral amino acids, zinc, estrogen and androgens, can contribute to the development of cancer, autoimmune disorders and viral infections (Afford and Randhawa, *Mol. Pathol.*, **2000**, 53, 55-63). Consequently, apoptosis is regulated under normal circumstances by the interaction of gene products that either induce or inhibit cell death and several gene products which modulate the apoptotic process have now been identified.

The most well-characterized apoptotic signaling cascade to date is that orchestrated by a family of cysteine proteases known as caspases. These enzymes activate apoptosis through proteolytic events triggered by one of several described mechanisms; including ligand binding to the cell surface death receptors of either the TNF or NGF receptor families, changes in mitochondrial integrity or chemical induction (Thornberry, *Br. Med. Bull.*, **1997**, 53, 478-490).

Caspases have been classified into two groups, initiator caspases and effector caspases, based upon their position in the apoptotic signaling pathway.

Initiator caspases include caspase 1, 2, 4, 5, 8, 9, 10 and 14 and these enzymes have the largest prodomains of all the caspase zymogens. These prodomains allow the initiator caspases to interact with other downstream substrates including other caspases. Initiator caspases are further divided into two groups based on their protein binding domains. Caspases 8 and 10 contain the DED (death effector domain) while caspases 1, 2, 4 and 9 contain the CARD (caspase recruitment domain) (Bratton et al., *Exp. Cell. Res.*, **2000**, 256, 27-33; Garcia-Calvo et al., *Cell. Death Differ.*, **1999**, 6, 362-369).

Effector caspases are activated by initiator caspases and include caspase 3, 6, 7, 11 and 13 and these contain a shorter prodomain. Once activated, effector caspases then cleave a number of structural and regulatory proteins within

the cell (Bratton et al., *Exp. Cell. Res.*, **2000**, 256, 27-33; Garcia-Calvo et al., *Cell. Death Differ.*, **1999**, 6, 362-369).

Caspase 7 (also known as CASP7, MCH3, ICE-LAP3 and CMH-1) is an effector caspase expressed in all tissues except brain (Duan et al., *J. Biol. Chem.*, **1996**, 271, 1621-1625; Fernandes-Alnemri et al., *Cancer Res.*, **1995**, 55, 6045-6052; Juan et al., *Genomics*, **1997**, 40, 86-93; Lippke et al., *J. Biol. Chem.*, **1996**, 271, 1825-1828). Characterization of the gene revealed that caspase 7 exists as two different isoforms generated by alternative splicing (Fernandes-Alnemri et al., *Cancer Res.*, **1995**, 55, 6045-6052).

Disclosed in the PCT publication are the polypeptide and nucleotide sequences of caspase 7 as are recombinant molecules encoding caspase 7, host cells transformed with said recombinant molecules and antibodies against the caspase 7 protein (Su and Lippke, **1997**).

Caspase 7 has been localized to human chromosome 10q25, a region which includes tumor suppressor genes believed to be associated with tumor progression in malignant gliomas and melanomas (Bullrich et al., *Genomics*, **1996**, 36, 362-365). In addition, increased activation of caspase 7 with concomitant induction of apoptosis has been noted in several cancers including prostate cancer and leukemias (King et al., *Leukemia*, **1998**, 12, 1553-1560; Marcelli et al., *Cancer Res.*, **1998**, 58, 76-83; Marcelli et al., *Cancer Res.*, **1999**, 59, 382-390).

Furthermore, it has been shown that caspase 7 affects other downstream pathways via cleavage of protein substrates involved in the stress response (Germain et al., *J. Biol. Chem.*, **1999**, 274, 28379-28384) and inflammation (Behrens et al., *FEBS Lett.*, **2000**, 466, 143-147).

Collectively, these data suggest that modulation of caspase 7 would render opportunity to treat patients with various cancers, inflammatory conditions and deregulated apoptotic pathologic conditions.

Several types of broad-spectrum caspase inhibitors have been identified for the treatment of deregulated bone

metabolism (Harada et al., 2000; Reszka, 1999), as immunomodulatory agents (Gunasekera et al., 2000) and as combination therapies for the regulation of blood cholesterol (Reszka, 1999). These non-specific caspase inhibitors fall  
5 into three main classes; peptide-based molecules that mimic caspase substrates, small molecules and naturally-occurring caspase inhibitors or decoys (Deveraux et al., *Embo J.*, 1998, 17, 2215-2223; Dong et al., *Biochem. J.*, 2000, 347 Pt 3, 669-677; Gunasekera et al., 2000; Harada et al., 2000; Reszka,  
10 1999; Reszka, 1999; Robidoux et al., 2000; Spruce et al., 1999).

Strategies aimed at modulating caspase 7 function have involved the use of antibodies and molecules that block upstream entities such as the death receptors and broad-  
15 spectrum caspase inhibitors. Bowen et al. demonstrated the role of caspase 7 in apoptosis using a 20-mer phosphodiester antisense oligonucleotide targeted to caspase 7. In this study it was shown that antisense depletion of caspase 7 slowed the onset of apoptosis in TSU-Prp1 prostate cancer  
20 cells. This antisense oligonucleotide, however, did not inhibit apoptosis to any greater extent than peptide inhibitors of cysteine proteases in general (Bowen et al., *Cell Death Differ.*, 1999, 6, 394-401).

Currently there exists a need to identify methods of  
25 modulating apoptosis for the therapeutic treatment of human diseases and it is believed that caspases modulators will be integral to these methods.

Antisense technology is emerging as an effective means for reducing the expression of specific gene products and may  
30 therefore prove to be uniquely useful in a number of therapeutic, diagnostic, and research applications for the modulation of gene expression and cellular processes.

There are no known therapeutic agents which effectively inhibit the synthesis of caspase 7 and the present invention  
35 provides compositions and methods for modulating caspase 7 expression, including modulation of aberrant forms of caspase 7, including alternatively spliced forms.

**SUMMARY OF THE INVENTION**

The present invention is directed to compounds, particularly antisense oligonucleotides, which are targeted to a nucleic acid encoding caspase 7, and which modulate the expression of caspase 7. Pharmaceutical and other compositions comprising the compounds of the invention are also provided. Further provided are methods of modulating the expression of caspase 7 in cells or tissues comprising contacting said cells or tissues with one or more of the antisense compounds or compositions of the invention. Further provided are methods of treating an animal, particularly a human, suspected of having or being prone to a disease or condition associated with expression of caspase 7 by administering a therapeutically or prophylactically effective amount of one or more of the antisense compounds or compositions of the invention.

**DETAILED DESCRIPTION OF THE INVENTION**

The present invention employs oligomeric compounds, particularly antisense oligonucleotides, for use in modulating the function of nucleic acid molecules encoding caspase 7, ultimately modulating the amount of caspase 7 produced. This is accomplished by providing antisense compounds which specifically hybridize with one or more nucleic acids encoding caspase 7. As used herein, the terms "target nucleic acid" and "nucleic acid encoding caspase 7" encompass DNA encoding caspase 7, RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of RNA



to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of caspase 7. In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a particular nucleic acid, in the context of this invention, is a multistep process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target is a nucleic acid molecule encoding caspase 7. The targeting process also includes determination of a site or sites within this gene for the antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present invention, a preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in

vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in

5 prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of  
10 the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used in vivo to initiate translation of an mRNA molecule transcribed from a gene encoding caspase 7, regardless of the sequence(s) of such codons.

15 It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and  
20 "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon  
25 region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which  
30 is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA  
35 in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding

nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation  
5 termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap  
10 structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as  
15 "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. mRNA splice sites, i.e., intron-exon junctions, may also be preferred target regions, and are  
20 particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. It has also been found that  
25 introns can also be effective, and therefore preferred, target regions for antisense compounds targeted, for example, to DNA or pre-mRNA.

Once one or more target sites have been identified, oligonucleotides are chosen which are sufficiently  
30 complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick, Hoogsteen or  
35 reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases. For example, adenine and thymine are complementary nucleobases which pair through the

formation of hydrogen bonds. "Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to each other at that position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that the sequence of an antisense compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. An antisense compound is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed.

Antisense and other compounds of the invention which hybridize to the target and inhibit expression of the target are identified through experimentation, and the sequences of these compounds are hereinbelow identified as preferred embodiments of the invention. The target sites to which these preferred sequences are complementary are hereinbelow referred to as "active sites" and are therefore preferred sites for targeting. Therefore another embodiment of the

invention encompasses compounds which hybridize to these active sites.

Antisense compounds are commonly used as research reagents and diagnostics. For example, antisense  
5 oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes. Antisense compounds are also used, for example, to distinguish between functions of various members of a  
10 biological pathway. Antisense modulation has, therefore, been harnessed for research use.

The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotides have been employed as therapeutic  
15 moieties in the treatment of disease states in animals and man. Antisense oligonucleotide drugs, including ribozymes, have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic  
20 modalities that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans.

In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of  
25 ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-occurring portions  
30 which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.  
35 While antisense oligonucleotides are a preferred form of antisense compound, the present invention comprehends other oligomeric antisense compounds, including but not limited to

oligonucleotide mimetics such as are described below. The antisense compounds in accordance with this invention preferably comprise from about 8 to about 50 nucleobases (i.e. from about 8 to about 50 linked nucleosides).

- 5 Particularly preferred antisense compounds are antisense oligonucleotides, even more preferably those comprising from about 12 to about 30 nucleobases. Antisense compounds include ribozymes, external guide sequence (EGS) oligonucleotides (oligozymes), and other short catalytic RNAs or catalytic  
10 oligonucleotides which hybridize to the target nucleic acid and modulate its expression.

- As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such  
15 heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to  
20 either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a  
25 circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5'  
30 phosphodiester linkage.

- Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having  
35 modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this

specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

5 Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates, 5'-alkylene phosphonates and chiral  
10 phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, selenophosphates and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs  
15 of these, and those having inverted polarity wherein one or more internucleotide linkages is a 3' to 3', 5' to 5' or 2' to 2' linkage. Preferred oligonucleotides having inverted polarity comprise a single 3' to 3' linkage at the 3'-most internucleotide linkage i.e. a single inverted nucleoside  
20 residue which may be abasic (the nucleobase is missing or has a hydroxyl group in place thereof). Various salts, mixed salts and free acid forms are also included.

Representative United States patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S.: 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,194,599; 5,565,555; 5,527,899; 5,721,218; 5,672,697 and 5,625,050, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

Preferred modified oligonucleotide backbones that do not  
35 include a phosphorus atom therein have backbones that are  
formed by short chain alkyl or cycloalkyl internucleoside  
linkages, mixed heteroatom and alkyl or cycloalkyl

internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; 5 sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; riboacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide 10 backbones; amide backbones; and others having mixed N, O, S and CH<sub>2</sub> component parts.

Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S.: 5,034,506; 5,166,315; 5,185,444; 15 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; 5,792,608; 5,646,269 and 20 5,677,439, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of 25 the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the 30 sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide 35 portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S.: 5,539,082; 5,714,331; and



5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., *Science*, **1991**, 254, 1497-1500.

Most preferred embodiments of the invention are  
5 oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular -CH<sub>2</sub>-NH-O-CH<sub>2</sub>-, -CH<sub>2</sub>-N(CH<sub>3</sub>)-O-CH<sub>2</sub>- [known as a methylene (methylimino) or MMI backbone], -CH<sub>2</sub>-O-N(CH<sub>3</sub>)-CH<sub>2</sub>-, -CH<sub>2</sub>-N(CH<sub>3</sub>)-N(CH<sub>3</sub>)-CH<sub>2</sub>- and -O-N(CH<sub>3</sub>)-CH<sub>2</sub>-CH<sub>2</sub>- [wherein the native  
10 phosphodiester backbone is represented as -O-P-O-CH<sub>2</sub>-] of the above referenced U.S. patent 5,489,677, and the amide backbones of the above referenced U.S. patent 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. patent  
15 5,034,506.

Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or  
20 O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C<sub>1</sub> to C<sub>10</sub> alkyl or C<sub>2</sub> to C<sub>10</sub> alkenyl and alkynyl. Particularly preferred are O[(CH<sub>2</sub>)<sub>n</sub>O]<sub>m</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>OCH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub>, O(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>ONH<sub>2</sub>, and O(CH<sub>2</sub>)<sub>n</sub>ON[(CH<sub>2</sub>)<sub>m</sub>CH<sub>3</sub>]<sub>2</sub>, where n and m are from 1 to about  
25 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C<sub>1</sub> to C<sub>10</sub> lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH<sub>3</sub>, OCN, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, SOCH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, heterocycloalkyl,  
30 heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and  
35 other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O-CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al.,

Helv. Chim. Acta, 1995, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylaminooxyethoxy, i.e., a  $O(CH_2)_2ON(CH_3)_2$  group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>2</sub>)<sub>2</sub>, also described in examples hereinbelow.

A further preferred modification includes Locked Nucleic Acids (LNAs) in which the 2'-hydroxyl group is linked to the 3' or 4' carbon atom of the sugar ring thereby forming a bicyclic sugar moiety. The linkage is preferably a methylene (-CH<sub>2</sub>)<sub>n</sub> group bridging the 2' oxygen atom and the 3' or 4' carbon atom wherein n is 1 or 2. LNAs and preparation thereof are described in WO 98/39352 and WO 99/14226.

Other preferred modifications include 2'-methoxy (2'-O-CH<sub>3</sub>), 2'-aminopropoxy (2'-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>), 2'-allyl (2'-CH<sub>2</sub>-CH=CH<sub>2</sub>), 2'-O-allyl (2'-O-CH<sub>2</sub>-CH=CH<sub>2</sub>) and 2'-fluoro (2'-F). The 2'-modification may be in the arabino (up) position or ribo (down) position. A preferred 2'-arabino modification is 2'-F. Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S.: 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; 5,792,747; and 5,700,920, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural"

nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl ( $-C\equiv C-CH_3$ ) uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 2-F-adenine, 2-amino-adenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further modified nucleobases include tricyclic pyrimidines such as phenoxazine cytidine(1H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-clamps such as a substituted phenoxazine cytidine (e.g. 9-(2-aminoethoxy)-H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido[4,5-b]indol-2-one), pyridoindole cytidine (H-pyrido[3',2':4,5]pyrrolo[2,3-d]pyrimidin-2-one). Modified nucleobases may also include those in which the purine or pyrimidine base is replaced with other heterocycles, for example 7-deaza-adenine, 7-deazaguanosine, 2-aminopyridine and 2-pyridone. Further nucleobases include those disclosed in United States Patent No. 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., *Angewandte Chemie*, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and*

Applications, pages 289-302, Crooke, S.T. and Lebleu, B., ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. 3,687,808, as well as U.S.: 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121; 5,596,091; 5,614,617; 5,645,985; 5,830,653; 5,763,588; 6,005,096; and 5,681,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference, and United States patent 5,750,692, which is commonly owned with the instant application and also herein incorporated by reference.

Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. The compounds of the invention can include conjugate groups covalently bound to functional groups such as primary or secondary hydroxyl groups. Conjugate groups of the invention include intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of

oligomers, and groups that enhance the pharmacokinetic properties of oligomers. Typical conjugates groups include cholesterol, lipids, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention, include groups that improve oligomer uptake, enhance oligomer resistance to degradation, and/or strengthen sequence-specific hybridization with RNA. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve oligomer uptake, distribution, metabolism or excretion. Representative conjugate groups are disclosed in International Patent Application PCT/US92/09196, filed October 23, 1992 the entire disclosure of which is incorporated herein by reference. Conjugate moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, **1989**, *86*, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, **1994**, *4*, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, **1992**, *660*, 306-309; Manoharan et al., *Bioorg. Med. Chem. Lett.*, **1993**, *3*, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, **1992**, *20*, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, **1991**, *10*, 1111-1118; Kabanov et al., *FEBS Lett.*, **1990**, *259*, 327-330; Svinarchuk et al., *Biochimie*, **1993**, *75*, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, **1995**, *36*, 3651-3654; Shea et al., *Nucl. Acids Res.*, **1990**, *18*, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, **1995**, *14*, 969-973), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, **1995**, *36*, 3651-3654), a palmityl moiety (Mishra et al., *Biochim. Biophys. Acta*, **1995**, *1264*, 229-237), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., *J.*

*Pharmacol. Exp. Ther.*, 1996, 277, 923-937. Oligonucleotides of the invention may also be conjugated to active drug substances, for example, aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-

- 5 pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folinic acid, a benzothiadiazide, chlorothiazide, a diazepine, indomethicin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic. Oligonucleotide-drug  
10 conjugates and their preparation are described in United States Patent Application 09/334,130 (filed June 15, 1999) which is incorporated herein by reference in its entirety.

- Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but  
15 are not limited to, U.S.: 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779;  
20 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667;  
25 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference.

- 30 It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an oligonucleotide. The present invention also includes  
35 antisense compounds which are chimeric compounds. "Chimeric" antisense compounds or "chimerae," in the context of this invention, are antisense compounds, particularly

oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional region of the oligonucleotide may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of oligonucleotide inhibition of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

Chimeric antisense compounds of the invention may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Such compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S.: 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

The antisense compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for

such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, CA). Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives.

The antisense compounds of the invention are synthesized in vitro and do not include antisense compositions of biological origin, or genetic vector constructs designed to direct the in vivo synthesis of antisense molecules. The compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption assisting formulations include, but are not limited to, U.S.: 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of which is herein incorporated by reference.

The antisense compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents.

The term "prodrug" indicates a therapeutic agent that is prepared in an inactive form that is converted to an active



form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. In particular, prodrug versions of the oligonucleotides of the invention are prepared as SATE  
5 [(S-acetyl-2-thioethyl) phosphat] derivatives according to the methods disclosed in WO 93/24510 to Gosselin et al., published December 9, 1993 or in WO 94/26764 and U.S. 5,770,713 to Imbach et al.

The term "pharmaceutically acceptable salts" refers to  
10 physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

Pharmaceutically acceptable base addition salts are  
15 formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chloroprocaine, choline,  
20 diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," *J. of Pharma Sci.*, 1977, 66, 1-19). The base addition salts of said acidic compounds are prepared by contacting the free acid form with a sufficient  
25 amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional manner. The free acid forms differ from their respective salt forms somewhat in certain  
30 physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention. As used herein, a "pharmaceutical addition salt" includes a pharmaceutically acceptable salt of an acid form of one of the components of  
35 the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and

phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example

5 hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric

10 acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or

15 isonicotinic acid; and with amino acids, such as the 20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic

20 acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid.

25 Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or

30 hydrogen carbonates are also possible.

For oligonucleotides, preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and

35 spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the

like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine.

10 The antisense compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an animal, preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of caspase 15 7 is treated by administering antisense compounds in accordance with this invention. The compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of an antisense compound to a suitable pharmaceutically acceptable diluent or carrier. Use 20 of the antisense compounds and methods of the invention may also be useful prophylactically, e.g., to prevent or delay infection, inflammation or tumor formation, for example.

The antisense compounds of the invention are useful for research and diagnostics, because these compounds hybridize 25 to nucleic acids encoding caspase 7, enabling sandwich and other assays to easily be constructed to exploit this fact. Hybridization of the antisense oligonucleotides of the invention with a nucleic acid encoding caspase 7 can be detected by means known in the art. Such means may include 30 conjugation of an enzyme to the oligonucleotide, radiolabelling of the oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of caspase 7 in a sample may also be prepared.

35 The present invention also includes pharmaceutical compositions and formulations which include the antisense compounds of the invention. The pharmaceutical compositions

of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration.

- Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful. Preferred topical formulations include those in which the oligonucleotides of the invention are in admixture with a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Preferred lipids and liposomes include neutral (e.g. dioleoylphosphatidyl DOPE ethanolamine, dimyristoylphosphatidyl choline DMPC, distearoylphosphatidyl choline) negative (e.g. dimyristoylphosphatidyl glycerol DMPG) and cationic (e.g. dioleoyltetramethylaminopropyl DOTAP and dioleoylphosphatidyl ethanolamine DOTMA). Oligonucleotides of the invention may be encapsulated within liposomes or may form complexes thereto, in particular to cationic liposomes. Alternatively, oligonucleotides may be complexed to lipids, in particular to cationic lipids. Preferred fatty acids and esters include but are not limited arachidonic acid, oleic acid, eicosanoic acid, lauric acid, caprylic acid, capric acid, myristic acid,

palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein, dilaurin, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a C<sub>1-10</sub> alkyl ester (e.g.

5 isopropylmyristate IPM), monoglyceride, diglyceride or pharmaceutically acceptable salt thereof. Topical formulations are described in detail in United States patent application 09/315,298 filed on May 20, 1999 which is incorporated herein by reference in its entirety.

10 Compositions and formulations for oral administration include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, capsules, gel capsules, sachets, tablets or minitables. Thickeners, flavoring agents, diluents, 15 emulsifiers, dispersing aids or binders may be desirable. Preferred oral formulations are those in which oligonucleotides of the invention are administered in conjunction with one or more penetration enhancers surfactants and chelators. Preferred surfactants include 20 fatty acids and/or esters or salts thereof, bile acids and/or salts thereof. Preferred bile acids/salts include chenodeoxycholic acid (CDCA) and ursodeoxychenodeoxycholic acid (UDCA), cholic acid, dehydrocholic acid, deoxycholic acid, gluccholic acid, glycholic acid, glycodeoxycholic acid, 25 taurocholic acid, taurodeoxycholic acid, sodium tauro-24,25-dihydro-fusidate, sodium glycodihydrofusidate,. Preferred fatty acids include arachidonic acid, undecanoic acid, oleic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, 30 dicaprate, tricaprate, monoolein, dilaurin, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a monoglyceride, a diglyceride or a pharmaceutically acceptable salt thereof (e.g. sodium). Also preferred are combinations of penetration enhancers, for 35 example, fatty acids/salts in combination with bile acids/salts. A particularly preferred combination is the

sodium salt of lauric acid, capric acid and UDCA. Further penetration enhancers include polyoxyethylene-9-lauryl ether, polyoxyethylene-20-cetyl ether. Oligonucleotides of the invention may be delivered orally in granular form including

5 sprayed dried particles, or complexed to form micro or nanoparticles. Oligonucleotide complexing agents include poly-amino acids; polyimines; polyacrylates; polyalkylacrylates, polyoxethanes, polyalkylcyanoacrylates; cationized gelatins, albumins, starches, acrylates,

10 polyethyleneglycols (PEG) and starches; polyalkylcyanoacrylates; DEAE-derivatized polyimines, pollulans, celluloses and starches. Particularly preferred complexing agents include chitosan, N-trimethylchitosan, poly-L-lysine, polyhistidine, polyornithine, polyspermines,

15 protamine, polyvinylpyridine, polythiodiethylamino-methylethylene P(TDAE), polyaminostyrene (e.g. p-amino), poly(methylcyanoacrylate), poly(ethylcyanoacrylate), poly(butylcyanoacrylate), poly(isobutylcyanoacrylate), poly(isohexylcyanoacrylate), DEAE-methacrylate, DEAE-

20 hexylacrylate, DEAE-acrylamide, DEAE-albumin and DEAE-dextran, polymethylacrylate, polyhexylacrylate, poly(D,L-lactic acid), poly(DL-lactic-co-glycolic acid (PLGA), alginate, and polyethyleneglycol (PEG). Oral formulations for oligonucleotides and their preparation are described in

25 detail in United States applications 08/886,829 (filed July 1, 1997), 09/108,673 (filed July 1, 1998), 09/256,515 (filed February 23, 1999), 09/082,624 (filed May 21, 1998) and 09/315,298 (filed May 20, 1999) each of which is incorporated herein by reference in their entirety.

30 Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and

35 other pharmaceutically acceptable carriers or excipients.

Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and

liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids.

5 The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active  
10 ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the  
15 product.

The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid syrups, soft gels, suppositories, and enemas. The  
20 compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or  
25 dextran. The suspension may also contain stabilizers.

In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams. Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams,  
30 jellies and liposomes. While basically similar in nature these formulations vary in the components and the consistency of the final product. The preparation of such compositions and formulations is generally known to those skilled in the pharmaceutical and formulation arts and may be applied to the  
35 formulation of the compositions of the present invention.

Emulsions

The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are typically heterogenous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1  $\mu\text{m}$  in diameter. (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 301). Emulsions are often biphasic systems comprising of two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions may be either water-in-oil (w/o) or of the oil-in-water (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to the dispersed phases and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a



system of oil droplets enclosed in globules of water stabilized in an oily continuous provides an o/w/o emulsion.

Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199). Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the preparation of formulations. Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet  
5 retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. These include polar inorganic solids, such as heavy metal  
10 hydroxides, nonswelling clays such as bentonite, attapulgite, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

A large variety of non-emulsifying materials are also  
15 included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker  
20 (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Hydrophilic colloids or hydrocolloids include naturally  
25 occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers (for example, carbomers, cellulose  
30 ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming strong interfacial films around the dispersed-phase droplets and by increasing the viscosity of the external phase.

35 Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may readily support the growth of microbes, these

formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of reasons of ease of formulation, efficacy from an absorption and bioavailability standpoint. (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

In one embodiment of the present invention, the compositions of oligonucleotides and nucleic acids are formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient

amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system.

Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, in: *Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215). Microemulsions commonly are prepared via a combination of three to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric packing of the polar heads and hydrocarbon tails of the surfactant molecules (Schott, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 271).

The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprate (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (SO750), decaglycerol

decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides (Constantinides et al., *Pharmaceutical Research*, **1994**, *11*, 1385-1390; Ritschel, *Meth. Find. Exp. Clin. Pharmacol.*, **1993**, *13*, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides et al., *Pharmaceutical Research*, **1994**, *11*, 1385; Ho et al., *J. Pharm. Sci.*, **1996**, *85*, 138-143). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be particularly advantageous when formulating thermolabile drugs, peptides or

oligonucleotides. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of oligonucleotides and nucleic acids from the gastrointestinal tract, as well as improve the local cellular uptake of oligonucleotides and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

Microemulsions of the present invention may also contain additional components and additives such as sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the oligonucleotides and nucleic acids of the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories - surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p. 92). Each of these classes has been discussed above.

#### Liposomes

There are many organized surfactant structures besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the

advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently with the cell wall, are taken up by macrophages *in vivo*.

In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through such fine pores.

Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes. As the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis.

Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang et al., *Biochem. Biophys. Res. Commun.*, **1987**, 147, 980-985).

Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou et al., *Journal of Controlled Release*, **1992**, 19, 269-274).

One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.



Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of  
5 interferon via other means (e.g. as a solution or as an emulsion) were ineffective (Weiner et al., *Journal of Drug Targeting*, **1992**, 2, 405-410). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon  
10 using an aqueous system, and concluded that the liposomal formulation was superior to aqueous administration (du Plessis et al., *Antiviral Research*, **1992**, 18, 259-265).

Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin,  
15 in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal formulations comprising Novasome™ I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome™ II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to  
20 deliver cyclosporin-A into the dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of cyclosporin-A into different layers of the skin (Hu et al. *S.T.P. Pharma. Sci.*, **1994**, 4, 6, 466).

Liposomes also include "sterically stabilized"  
liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation  
lifetimes relative to liposomes lacking such specialized  
30 lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside GM<sub>1</sub>, or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol  
35 (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least

for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., *FEBS Letters*, **1987**, 223, 42; Wu et al., *Cancer Research*, **1993**, 53, 3765).

Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (*Ann. N.Y. Acad. Sci.*, **1987**, 507, 64) reported the ability of monosialoganglioside GM<sub>1</sub>, galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (*Proc. Natl. Acad. Sci. U.S.A.*, **1988**, 85, 6949). U.S. Patent No. 4,837,028 and WO 88/04924, both to Allen et al., disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside GM<sub>1</sub> or a galactocerebroside sulfate ester. U.S. Patent No. 5,543,152 (Webb et al.) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-sn-dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim et al.).

Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (*Bull. Chem. Soc. Jpn.*, **1980**, 53, 2778) described liposomes comprising a nonionic detergent, 2C<sub>12</sub>15G, that contains a PEG moiety. Illum et al. (*FEBS Lett.*, **1984**, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Patent Nos. 4,426,330 and 4,534,899). Klibanov et al. (*FEBS Lett.*, **1990**, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume et al. (*Biochimica et Biophysica Acta*, **1990**, 1029, 91) extended such observations to other PEG-

derivatized phospholipids, e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle et al. (U.S. Patent Nos. 5,013,556 and 5,356,633) and Martin et al. (U.S. Patent No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Patent No. 5,225,212 (both to Martin et al.) and in WO 94/20073 (Zalipsky et al.) Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi et al.). U.S. Patent Nos. 5,540,935 (Miyazaki et al.) and 5,556,948 (Tagawa et al.) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry et al. discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Patent No. 5,264,221 to Tagawa et al. discloses protein-bonded liposomes and asserts that the contents of such liposomes may include an antisense RNA. U.S. Patent No. 5,665,710 to Rahman et al. describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787 to Love et al. discloses liposomes comprising antisense oligonucleotides targeted to the raf gene.

Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, e.g. they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without

fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and

sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

5 If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

10 If the surfactant molecule has the ability to carry either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

15 The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

#### 20 Penetration Enhancers

In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Most drugs are present in solution in both ionized  
25 and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of  
30 non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-  
35 surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92). Each of the above mentioned

classes of penetration enhancers are described below in greater detail.

Surfactants: In connection with the present invention, surfactants (or "surface-active agents") are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, with the result that absorption of oligonucleotides through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92); and perfluorochemical emulsions, such as FC-43. Takahashi et al., *J. Pharm. Pharmacol.*, 1988, 40, 252).

Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein (1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C<sub>10-18</sub> alkyl esters thereof (e.g., methyl, isopropyl and t-butyl), and mono- and di-glycerides thereof (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; El Hariri et al., *J. Pharm. Pharmacol.*, 1992, 44, 651-654).

Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York, 1996, pp. 934-935).

Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. The bile salts of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucolic acid (sodium glucolate), glycholic acid (sodium glycocholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydro-fusidate (STDHF), sodium glycodihydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Swinyard, Chapter 39 In: *Remington's Pharmaceutical Sciences*, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, 1990, pages 782-783; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Yamamoto et al., *J. Pharm. Exp. Ther.*, 1992, 263, 25; Yamashita et al., *J. Pharm. Sci.*, 1990, 79, 579-583).

Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that absorption of oligonucleotides through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal ion for catalysis and are thus inhibited by chelating agents (Jarrett, *J. Chromatogr.*, 1993, 618, 315-339). Chelating agents of the invention include but are not limited to disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and

homovanilate), N-acyl derivatives of collagen, laureth-9 and N-amino acyl derivatives of beta-diketones (enamines) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, **1991**, page 92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, **1990**, 7, 1-33; Buur et al., *J. Control Rel.*, **1990**, 14, 43-51).

Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as chelating agents or as surfactants but that nonetheless enhance absorption of oligonucleotides through the alimentary mucosa (Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, **1990**, 7, 1-33). This class of penetration enhancers include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, **1991**, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., *J. Pharm. Pharmacol.*, **1987**, 39, 621-626).

Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. Patent No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo et al., PCT Application WO 97/30731), are also known to enhance the cellular uptake of oligonucleotides.

Other agents may be utilized to enhance the penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

### 35 Carriers

Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used



herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which is inert (i.e., does not possess biological activity *per se*) but is recognized as a nucleic acid by *in vivo* processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioate oligonucleotide in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'-isothiocyano-stilbene-2,2'-disulfonic acid (Miyao et al., *Antisense Res. Dev.*, **1995**, 5, 115-121; Takakura et al., *Antisense & Nucl. Acid Drug Dev.*, **1996**, 6, 177-183).

#### Excipients

In contrast to a carrier compound, a "pharmaceutical carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (e.g.,

5 magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrants (e.g., starch, sodium starch glycolate, etc.); and wetting agents (e.g., sodium lauryl sulphate, etc.).

10 Pharmaceutically acceptable organic or inorganic excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin,

15 hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-aqueous solutions in common solvents such as alcohols, or solutions of the nucleic acids in liquid or solid oil bases.

20 The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration which do not deleteriously react with nucleic acids can be used.

25 Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

30

#### Other Components

The compositions of the present invention may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their

35 art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example,

antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes,

5 flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and,  
10 if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s) of the  
15 formulation.

Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

20 Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more antisense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents include  
25 but are not limited to daunorubicin, daunomycin, dactinomycin, doxorubicin, epirubicin, idarubicin, esorubicin, bleomycin, mafosfamide, ifosfamide, cytosine arabinoside, bis-chloroethylnitrosurea, busulfan, mitomycin C, actinomycin D, mithramycin, prednisone,  
30 hydroxyprogesterone, testosterone, tamoxifen, dacarbazine, procarbazine, hexamethylmelamine, pentamethylmelamine, mitoxantrone, amsacrine, chlorambucil, methylcyclohexylnitrosurea, nitrogen mustards, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine,  
35 cytarabine, 5-azacytidine, hydroxyurea, deoxycoformycin, 4-hydroxyperoxycyclophosphoramide, 5-fluorouracil (5-FU), 5-fluorodeoxyuridine (5-FUDR), methotrexate (MTX), colchicine,

taxol, vincristine, vinblastine, etoposide (VP-16), trimetrexate, irinotecan, topotecan, gemcitabine, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed. 1987, pp.

5 1206-1228, Berkow et al., eds., Rahway, N.J. When used with the compounds of the invention, such chemotherapeutic agents may be used individually (e.g., 5-FU and oligonucleotide), sequentially (e.g., 5-FU and oligonucleotide for a period of time followed by MTX and oligonucleotide), or in combination  
10 with one or more other such chemotherapeutic agents (e.g., 5-FU, MTX and oligonucleotide, or 5-FU, radiotherapy and oligonucleotide). Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not  
15 limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-antisense  
20 chemotherapeutic agents are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

In another related embodiment, compositions of the invention may contain one or more antisense compounds,  
25 particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted to a second nucleic acid target. Numerous examples of antisense compounds are known in the art. Two or more combined compounds may be used together or sequentially.

30 The formulation of therapeutic compositions and their subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several  
35 months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body

of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on EC<sub>50</sub>s found to be effective in in vitro and in vivo animal models. In general, dosage is from 0.01 ug to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 ug to 100 g per kg of body weight, once or more daily, to once every 20 years.

While the present invention has been described with specificity in accordance with certain of its preferred embodiments, the following examples serve only to illustrate the invention and are not intended to limit the same.

## EXAMPLES

## Example 1

5 Nucleoside Phosphoramidites for Oligonucleotide Synthesis  
Deoxy and 2'-alkoxy amidites

2'-Deoxy and 2'-methoxy beta-cyanoethyl-diisopropyl  
phosphoramidites were purchased from commercial sources (e.g.  
ChemGenes, Needham MA or Glen Research, Inc. Sterling VA).  
10 Other 2'-O-alkoxy substituted nucleoside amidites are  
prepared as described in U.S. Patent 5,506,351, herein  
incorporated by reference. For oligonucleotides synthesized  
using 2'-alkoxy amidites, the standard cycle for unmodified  
oligonucleotides was utilized, except the wait step after  
15 pulse delivery of tetrazole and base was increased to 360  
seconds.

Oligonucleotides containing 5-methyl-2'-deoxycytidine  
(5-Me-C) nucleotides were synthesized according to published  
methods [Sanghvi, et. al., *Nucleic Acids Research*, **1993**, 21,  
20 3197-3203] using commercially available phosphoramidites  
(Glen Research, Sterling VA or ChemGenes, Needham MA).

## 2'-Fluoro amidites

## 2'-Fluorodeoxyadenosine amidites

25 2'-fluoro oligonucleotides were synthesized as described  
previously [Kawasaki, et. al., *J. Med. Chem.*, **1993**, 36, 831-  
841] and United States patent 5,670,633, herein incorporated  
by reference. Briefly, the protected nucleoside N6-benzoyl-  
2'-deoxy-2'-fluoroadenosine was synthesized utilizing  
30 commercially available 9-beta-D-arabinofuranosyladenine as  
starting material and by modifying literature procedures  
whereby the 2'-alpha-fluoro atom is introduced by a S<sub>N</sub>2-  
displacement of a 2'-beta-trityl group. Thus N6-benzoyl-9-  
beta-D-arabinofuranosyladenine was selectively protected in  
35 moderate yield as the 3',5'-ditetrahydropyranyl (THP)  
intermediate. Deprotection of the THP and N6-benzoyl groups  
was accomplished using standard methodologies and standard

methods were used to obtain the 5'-dimethoxytrityl- (DMT) and 5'-DMT-3'-phosphoramidite intermediates.

### 2'-Fluorodeoxyguanosine

5 The synthesis of 2'-deoxy-2'-fluoroguanosine was accomplished using tetraisopropylidisiloxanyl (TPDS) protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutryl-arabinofuranosylguanosine. Deprotection of the TPDS group  
10 was followed by protection of the hydroxyl group with THP to give diisobutryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation was followed by treatment of the crude product with fluoride, then deprotection of the THP groups. Standard methodologies were  
15 used to obtain the 5'-DMT- and 5'-DMT-3'-phosphoramidites.

### 2'-Fluorouridine

Synthesis of 2'-deoxy-2'-fluorouridine was accomplished by the modification of a literature procedure in which 2,2'-  
20 anhydro-1-beta-D-arabinofuranosyluracil was treated with 70% hydrogen fluoride-pyridine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

### 2'-Fluorodeoxycytidine

25 2'-deoxy-2'-fluorocytidine was synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N4-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

### 2'-O-(2-Methoxyethyl) modified amidites

2'-O-Methoxyethyl-substituted nucleoside amidites are prepared as follows, or alternatively, as per the methods of Martin, P., *Helvetica Chimica Acta*, 1995, 78, 486-504.

35 2,2'-Anhydro[1-(beta-D-arabinofuranosyl)-5-methyluridine]

5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenyl-carbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL). The mixture was heated to reflux, with stirring, allowing the evolved carbon dioxide to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The product formed a gum. The ether was decanted and the residue was dissolved in a minimum amount of methanol (ca. 400 mL). The solution was poured into fresh ether (2.5 L) to yield a stiff gum. The ether was decanted and the gum was dried in a vacuum oven (60°C at 1 mm Hg for 24 h) to give a solid that was crushed to a light tan powder (57 g, 85% crude yield). The NMR spectrum was consistent with the structure, contaminated with phenol as its sodium salt (ca. 5%). The material was used as is for further reactions (or it can be purified further by column chromatography using a gradient of methanol in ethyl acetate (10-25%) to give a white solid, mp 222-4°C).

#### 2'-O-Methoxyethyl-5-methyluridine

2,2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160°C. After heating for 48 hours at 155-160°C, the vessel was opened and the solution evaporated to dryness and triturated with MeOH (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate evaporated. The residue (280 g) was dissolved in CH<sub>3</sub>CN (600 mL) and evaporated. A silica gel column (3 kg) was packed in CH<sub>2</sub>Cl<sub>2</sub>/acetone/MeOH (20:5:3) containing 0.5% Et<sub>3</sub>NH. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with the packing solvent to give 160 g (63%) of



product. Additional material was obtained by reworking impure fractions.

**2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine**

5 2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue dissolved in pyridine (1.3 L). A first aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the mixture stirred at room temperature for one hour. A second  
10 aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the reaction stirred for an additional one hour. Methanol (170 mL) was then added to stop the reaction. HPLC showed the presence of approximately 70% product. The solvent was evaporated and triturated with CH<sub>3</sub>CN (200 mL).  
15 The residue was dissolved in CHCl<sub>3</sub> (1.5 L) and extracted with 2x500 mL of saturated NaHCO<sub>3</sub> and 2x500 mL of saturated NaCl. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated. 275 g of residue was obtained. The residue was purified on a 3.5 kg silica gel column, packed and eluted  
20 with EtOAc/hexane/acetone (5:5:1) containing 0.5% Et<sub>3</sub>NH. The pure fractions were evaporated to give 164 g of product. Approximately 20 g additional was obtained from the impure fractions to give a total yield of 183 g (57%).

25 **3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine**

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (106 g, 0.167 M), DMF/pyridine (750 mL of a 3:1 mixture prepared from 562 mL of DMF and 188 mL of pyridine) and  
30 acetic anhydride (24.38 mL, 0.258 M) were combined and stirred at room temperature for 24 hours. The reaction was monitored by TLC by first quenching the TLC sample with the addition of MeOH. Upon completion of the reaction, as judged by TLC, MeOH (50 mL) was added and the mixture evaporated at  
35 35°C. The residue was dissolved in CHCl<sub>3</sub> (800 mL) and extracted with 2x200 mL of saturated sodium bicarbonate and

2x200 mL of saturated NaCl. The water layers were back extracted with 200 mL of CHCl<sub>3</sub>. The combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product). The residue was purified on a 5 3.5 kg silica gel column and eluted using EtOAc/hexane(4:1). Pure product fractions were evaporated to yield 96 g (84%). An additional 1.5 g was recovered from later fractions.

**3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine**

10

A first solution was prepared by dissolving 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (96 g, 0.144 M) in CH<sub>3</sub>CN (700 mL) and set aside. Triethylamine (189 mL, 1.44 M) was added to a solution of triazole (90 g, 1.3 M) 15 in CH<sub>3</sub>CN (1 L), cooled to -5°C and stirred for 0.5 h using an overhead stirrer. POCl<sub>3</sub> was added dropwise, over a 30 minute period, to the stirred solution maintained at 0-10°C, and the resulting mixture stirred for an additional 2 hours. The first solution was added dropwise, over a 45 minute period, 20 to the latter solution. The resulting reaction mixture was stored overnight in a cold room. Salts were filtered from the reaction mixture and the solution was evaporated. The residue was dissolved in EtOAc (1 L) and the insoluble solids were removed by filtration. The filtrate was washed with 25 1x300 mL of NaHCO<sub>3</sub> and 2x300 mL of saturated NaCl, dried over sodium sulfate and evaporated. The residue was triturated with EtOAc to give the title compound.

**2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

30

A solution of 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141 M) in dioxane (500 mL) and NH<sub>4</sub>OH (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was evaporated and the residue azeotroped with MeOH (2x200 mL). The residue 35 was dissolved in MeOH (300 mL) and transferred to a 2 liter stainless steel pressure vessel. MeOH (400 mL) saturated

with NH<sub>3</sub> gas was added and the vessel heated to 100°C for 2 hours (TLC showed complete conversion). The vessel contents were evaporated to dryness and the residue was dissolved in EtOAc (500 mL) and washed once with saturated NaCl (200 mL).  
5 The organics were dried over sodium sulfate and the solvent was evaporated to give 85 g (95%) of the title compound.

**N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

10 2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) was added with stirring. After stirring for 3 hours, TLC showed the reaction to be approximately 95% complete. The solvent was evaporated and  
15 the residue azeotroped with MeOH (200 mL). The residue was dissolved in CHCl<sub>3</sub> (700 mL) and extracted with saturated NaHCO<sub>3</sub> (2x300 mL) and saturated NaCl (2x300 mL), dried over MgSO<sub>4</sub> and evaporated to give a residue (96 g). The residue was chromatographed on a 1.5 kg silica column using  
20 EtOAc/hexane (1:1) containing 0.5% Et<sub>3</sub>NH as the eluting solvent. The pure product fractions were evaporated to give 90 g (90%) of the title compound.

**N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine-3'-amidite**

25 N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (74 g, 0.10 M) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1 L). Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxy-tetra-(isopropyl)phosphite (40.5 mL, 0.123 M) were added with  
30 stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (TLC showed the reaction to be 95% complete). The reaction mixture was extracted with saturated NaHCO<sub>3</sub> (1x300 mL) and saturated NaCl (3x300 mL). The aqueous washes were back-extracted with  
35 CH<sub>2</sub>Cl<sub>2</sub> (300 mL), and the extracts were combined, dried over MgSO<sub>4</sub> and concentrated. The residue obtained was

chromatographed on a 1.5 kg silica column using EtOAc/hexane (3:1) as the eluting solvent. The pure fractions were combined to give 90.6 g (87%) of the title compound.

5    **2'-O-(Aminooxyethyl) nucleoside amidites and 2'-O-(dimethylaminooxyethyl) nucleoside amidites**

**2'-(Dimethylaminooxyethoxy) nucleoside amidites**

          2'-(Dimethylaminooxyethoxy) nucleoside amidites [also  
10    known in the art as 2'-O-(dimethylaminooxyethyl) nucleoside  
amidites] are prepared as described in the following  
paragraphs. Adenosine, cytidine and guanosine nucleoside  
amidites are prepared similarly to the thymidine (5-  
methyluridine) except the exocyclic amines are protected with  
15    a benzoyl moiety in the case of adenosine and cytidine and  
with isobutyryl in the case of guanosine.

**5'-O-tert-Butyldiphenylsilyl-O<sup>2</sup>-2'-anhydro-5-methyluridine**

20    O<sup>2</sup>-2'-anhydro-5-methyluridine (Pro. Bio. Sint., Varese, Italy, 100.0g, 0.416 mmol), dimethylaminopyridine (0.66g, 0.013eq, 0.0054mmol) were dissolved in dry pyridine (500 ml) at ambient temperature under an argon atmosphere and with mechanical stirring. tert-Butyldiphenylchlorosilane (125.8g,  
25    119.0mL, 1.1eq, 0.458mmol) was added in one portion. The reaction was stirred for 16 h at ambient temperature. TLC (R<sub>f</sub> 0.22, ethyl acetate) indicated a complete reaction. The solution was concentrated under reduced pressure to a thick oil. This was partitioned between dichloromethane (1 L) and  
30    saturated sodium bicarbonate (2x1 L) and brine (1 L). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to a thick oil. The oil was dissolved in a 1:1 mixture of ethyl acetate and ethyl ether (600mL) and the solution was cooled to  
35    -10°C. The resulting crystalline product was collected by filtration, washed with ethyl ether (3x200 mL) and dried

(40°C, 1mm Hg, 24 h) to 149g (74.8%) of white solid. TLC and NMR were consistent with pure product.

**5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine**

In a 2 L stainless steel, unstirred pressure reactor was added borane in tetrahydrofuran (1.0 M, 2.0 eq, 622 mL). In the fume hood and with manual stirring, ethylene glycol (350 mL, excess) was added cautiously at first until the evolution of hydrogen gas subsided. 5'-O-tert-Butyldiphenylsilyl-O<sup>2</sup>-2'-anhydro-5-methyluridine (149 g, 0.311 mol) and sodium bicarbonate (0.074 g, 0.003 eq) were added with manual stirring. The reactor was sealed and heated in an oil bath until an internal temperature of 160 °C was reached and then maintained for 16 h (pressure < 100 psig). The reaction vessel was cooled to ambient and opened. TLC (Rf 0.67 for desired product and Rf 0.82 for ara-T side product, ethyl acetate) indicated about 70% conversion to the product. In order to avoid additional side product formation, the reaction was stopped, concentrated under reduced pressure (10 to 1mm Hg) in a warm water bath (40-100°C) with the more extreme conditions used to remove the ethylene glycol. [Alternatively, once the low boiling solvent is gone, the remaining solution can be partitioned between ethyl acetate and water. The product will be in the organic phase.] The residue was purified by column chromatography (2kg silica gel, ethyl acetate-hexanes gradient 1:1 to 4:1). The appropriate fractions were combined, stripped and dried to product as a white crisp foam (84g, 50%), contaminated starting material (17.4g) and pure reusable starting material 20g. The yield based on starting material less pure recovered starting material was 58%. TLC and NMR were consistent with 99% pure product.

**2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-methyluridine**

5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine (20g, 36.98mmol) was mixed with triphenylphosphine (11.63g, 44.36mmol) and N-hydroxyphthalimide (7.24g, 44.36mmol). It was then dried over P<sub>2</sub>O<sub>5</sub> under high vacuum for two days at 40°C. The reaction mixture was flushed with argon and dry THF (369.8mL, Aldrich, sure seal bottle) was added to get a clear solution. Diethyl-azodicarboxylate (6.98mL, 44.36mmol) was added dropwise to the reaction mixture. The rate of addition is maintained such that resulting deep red coloration is just discharged before adding the next drop. After the addition was complete, the reaction was stirred for 4 hrs. By that time TLC showed the completion of the reaction (ethylacetate:hexane, 60:40). The solvent was evaporated in vacuum. Residue obtained was placed on a flash column and eluted with ethyl acetate:hexane (60:40), to get 2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-methyluridine as white foam (21.819 g, 86%).

20 **5'-O-tert-butyldiphenylsilyl-2'-O-([2-formadoximinooxy)ethyl]-5-methyluridine**

2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-methyluridine (3.1g, 4.5mmol) was dissolved in dry CH<sub>2</sub>Cl<sub>2</sub> (4.5mL) and methylhydrazine (300mL, 4.64mmol) was added dropwise at -10°C to 0°C. After 1 h the mixture was filtered, the filtrate was washed with ice cold CH<sub>2</sub>Cl<sub>2</sub> and the combined organic phase was washed with water, brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The solution was concentrated to get 2'-O-(aminooxyethyl) thymidine, which was then dissolved in MeOH (67.5mL). To this formaldehyde (20% aqueous solution, w/w, 1.1 eq.) was added and the resulting mixture was stirred for 1 h. Solvent was removed under vacuum; residue chromatographed to get 5'-O-tert-butyldiphenylsilyl-2'-O-([2-formadoximinooxy)ethyl]-5-methyluridine as white foam (1.95 g, 78%).

**5'-O-tert-Butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine**

5'-O-tert-butyldiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine (1.77g, 3.12mmol) was dissolved in a solution of 1M pyridinium p-toluenesulfonate (PPTS) in dry MeOH (30.6mL). Sodium cyanoborohydride (0.39g, 6.13mmol) was added to this solution at 10°C under inert atmosphere. The reaction mixture was stirred for 10 minutes at 10°C. After that the reaction vessel was removed from the ice bath and stirred at room temperature for 2 h, the reaction monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>). Aqueous NaHCO<sub>3</sub> solution (5%, 10mL) was added and extracted with ethyl acetate (2x20mL). Ethyl acetate phase was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, evaporated to dryness. Residue was dissolved in a solution of 1M PPTS in MeOH (30.6mL). Formaldehyde (20% w/w, 30mL, 3.37mmol) was added and the reaction mixture was stirred at room temperature for 10 minutes. Reaction mixture cooled to 10°C in an ice bath, sodium cyanoborohydride (0.39g, 6.13mmol) was added and reaction mixture stirred at 10°C for 10 minutes. After 10 minutes, the reaction mixture was removed from the ice bath and stirred at room temperature for 2 hrs. To the reaction mixture 5% NaHCO<sub>3</sub> (25mL) solution was added and extracted with ethyl acetate (2x25mL). Ethyl acetate layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness. The residue obtained was purified by flash column chromatography and eluted with 5% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to get 5'-O-tert-butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine as a white foam (14.6g, 80%).

**2'-O-(dimethylaminoxyethyl)-5-methyluridine**

Triethylamine trihydrofluoride (3.91mL, 24.0mmol) was dissolved in dry THF and triethylamine (1.67mL, 12mmol, dry, kept over KOH). This mixture of triethylamine-2HF was then added to 5'-O-tert-butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine (1.40g, 2.4mmol) and

stirred at room temperature for 24 hrs. Reaction was monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>). Solvent was removed under vacuum and the residue placed on a flash column and eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to get 2'-O-

5 (dimethylaminooxyethyl)-5-methyluridine (766mg, 92.5%).

5'-O-DMT-2'-O-(dimethylaminooxyethyl)-5-methyluridine  
2'-O-(dimethylaminooxyethyl)-5-methyluridine (750mg,  
2.17mmol) was dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight at  
10 40°C. It was then co-evaporated with anhydrous pyridine  
(20mL). The residue obtained was dissolved in pyridine  
(11mL) under argon atmosphere. 4-dimethylaminopyridine  
(26.5mg, 2.60mmol), 4,4'-dimethoxytrityl chloride (880mg,  
2.60mmol) was added to the mixture and the reaction mixture  
15 was stirred at room temperature until all of the starting  
material disappeared. Pyridine was removed under vacuum and  
the residue chromatographed and eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub>  
(containing a few drops of pyridine) to get 5'-O-DMT-2'-O-  
(dimethylamino-oxyethyl)-5-methyluridine (1.13g, 80%).

20

5'-O-DMT-2'-O-(2-N,N-dimethylaminooxyethyl)-5-  
methyluridine-3'-[(2-cyanoethyl)-N,N-  
diisopropylphosphoramidite]

5'-O-DMT-2'-O-(dimethylaminooxyethyl)-5-methyluridine  
25 (1.08g, 1.67mmol) was co-evaporated with toluene (20mL). To  
the residue N,N-diisopropylamine tetrazonide (0.29g,  
1.67mmol) was added and dried over P<sub>2</sub>O<sub>5</sub> under high vacuum  
overnight at 40°C. Then the reaction mixture was dissolved in  
anhydrous acetonitrile (8.4mL) and 2-cyanoethyl-N,N,N<sup>1</sup>,N<sup>1</sup>-  
30 tetraisopropylphosphoramidite (2.12mL, 6.08mmol) was added.  
The reaction mixture was stirred at ambient temperature for 4  
hrs under inert atmosphere. The progress of the reaction was  
monitored by TLC (hexane:ethyl acetate 1:1). The solvent was  
evaporated, then the residue was dissolved in ethyl acetate  
35 (70mL) and washed with 5% aqueous NaHCO<sub>3</sub> (40mL). Ethyl  
acetate layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and



concentrated. Residue obtained was chromatographed (ethyl acetate as eluent) to get 5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite] as a foam (1.04g, 74.9%).

5

**2'-(Aminooxyethoxy) nucleoside amidites**

2'-(Aminooxyethoxy) nucleoside amidites [also known in the art as 2'-O-(aminooxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, 10 cytidine and thymidine nucleoside amidites are prepared similarly.

**N2-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]**

15 The 2'-O-aminooxyethyl guanosine analog may be obtained by selective 2'-O-alkylation of diaminopurine riboside. Multigram quantities of diaminopurine riboside may be purchased from Schering AG (Berlin) to provide 2'-O-(2-ethylacetyl) diaminopurine riboside along with a minor amount 20 of the 3'-O-isomer. 2'-O-(2-ethylacetyl) diaminopurine riboside may be resolved and converted to 2'-O-(2-ethylacetyl)guanosine by treatment with adenosine deaminase. (McGee, D. P. C., Cook, P. D., Guinosso, C. J., WO 94/02501 25 A1 940203.) Standard protection procedures should afford 2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine and 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine which may be reduced to provide 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine. As before 30 the hydroxyl group may be displaced by N-hydroxyphthalimide via a Mitsunobu reaction, and the protected nucleoside may phosphitylated as usual to yield 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]. 35

**2'-dimethylaminoethoxyethoxy (2'-DMAEOE) nucleoside amidites**

2'-dimethylaminoethoxyethoxy nucleoside amidites (also known in the art as 2'-O-dimethylaminoethoxyethyl, i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>3</sub>)<sub>2</sub>, or 2'-DMAEOE nucleoside amidites) are prepared as follows. Other nucleoside amidites are prepared similarly.

**2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

2[2-(Dimethylamino)ethoxy]ethanol (Aldrich, 6.66 g, 50 mmol) is slowly added to a solution of borane in tetrahydrofuran (1 M, 10 mL, 10 mmol) with stirring in a 100 mL bomb. Hydrogen gas evolves as the solid dissolves. O<sup>2</sup>-,2'-anhydro-5-methyluridine (1.2 g, 5 mmol), and sodium bicarbonate (2.5 mg) are added and the bomb is sealed, placed in an oil bath and heated to 155°C for 26 hours. The bomb is cooled to room temperature and opened. The crude solution is concentrated and the residue partitioned between water (200 mL) and hexanes (200 mL). The excess phenol is extracted into the hexane layer. The aqueous layer is extracted with ethyl acetate (3x200 mL) and the combined organic layers are washed once with water, dried over anhydrous sodium sulfate and concentrated. The residue is columned on silica gel using methanol/methylene chloride 1:20 (which has 2% triethylamine) as the eluent. As the column fractions are concentrated a colorless solid forms which is collected to give the title compound as a white solid.

**5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

To 0.5 g (1.3 mmol) of 2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine in anhydrous pyridine (8 mL), triethylamine (0.36 mL) and dimethoxytrityl chloride (DMT-Cl, 0.87 g, 2 eq.) are added and stirred for 1 hour. The reaction mixture is poured into water (200 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (2x200 mL). The combined CH<sub>2</sub>Cl<sub>2</sub> layers are washed with saturated NaHCO<sub>3</sub> solution, followed by saturated NaCl

solution and dried over anhydrous sodium sulfate. Evaporation of the solvent followed by silica gel chromatography using MeOH:CH<sub>2</sub>Cl<sub>2</sub>:Et<sub>3</sub>N (20:1, v/v, with 1% triethylamine) gives the title compound.

5

**5'-O-Dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)-ethyl]-5-methyl uridine-3'-O-(cyanoethyl-N,N-diisopropyl)phosphoramidite**

Diisopropylaminotetrazolide (0.6 g) and 2-cyanoethoxy-  
10 N,N-diisopropyl phosphoramidite (1.1 mL, 2 eq.) are added to a solution of 5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyluridine (2.17 g, 3 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) under an atmosphere of argon. The reaction mixture is stirred overnight and the solvent  
15 evaporated. The resulting residue is purified by silica gel flash column chromatography with ethyl acetate as the eluent to give the title compound.

**Example 2**

**20 Oligonucleotide synthesis**

Unsubstituted and substituted phosphodiester (P=O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine.

25 Phosphorothioates (P=S) are synthesized as for the phosphodiester oligonucleotides except the standard oxidation bottle was replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation  
30 wait step was increased to 68 sec and was followed by the capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55°C (18 h), the oligonucleotides were purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M NaCl solution.

35 Phosphinate oligonucleotides are prepared as described in U.S. Patent 5,508,270, herein incorporated by reference.

Alkyl phosphonate oligonucleotides are prepared as described in U.S. Patent 4,469,863, herein incorporated by reference.

3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Patents 5,610,289 or 5,625,050, herein incorporated by reference.

Phosphoramidite oligonucleotides are prepared as described in U.S. Patent, 5,256,775 or U.S. Patent 5,366,878, herein incorporated by reference.

Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference.

3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Patent 5,476,925, herein incorporated by reference.

Phosphotriester oligonucleotides are prepared as described in U.S. Patent 5,023,243, herein incorporated by reference.

Borano phosphate oligonucleotides are prepared as described in U.S. Patents 5,130,302 and 5,177,198, both herein incorporated by reference.

### Example 3

#### Oligonucleoside Synthesis

Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedimethylhydrazo linked oligonucleosides, also identified as MDH linked oligonucleosides, and methylenecarbonylamino linked oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligonucleosides, also identified as amide-4 linked oligonucleosides, as well as mixed backbone compounds having, for instance, alternating MMI and P=O or P=S linkages are prepared as described in U.S. Patents 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289, all of which are herein incorporated by reference.

Formacetal and thioformacetal linked oligonucleosides are prepared as described in U.S. Patents 5,264,562 and 5,264,564, herein incorporated by reference.

Ethylene oxide linked oligonucleosides are prepared as described in U.S. Patent 5,223,618, herein incorporated by reference.

#### **Example 4**

##### **PNA Synthesis**

Peptide nucleic acids (PNAs) are prepared in accordance with any of the various procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, *Bioorganic & Medicinal Chemistry*, 1996, 4, 5-23. They may also be prepared in accordance with U.S. Patents 5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

#### **Example 5**

##### **Synthesis of Chimeric Oligonucleotides**

Chimeric oligonucleotides, oligonucleosides or mixed oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3' "wing" segments of linked nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers".

##### **[2'-O-Me]--[2'-deoxy]--[2'-O-Me] Chimeric Phosphorothioate Oligonucleotides**

Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligonucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 380B, as above. Oligonucleotides are synthesized using the automated

synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphor-  
amidite for the DNA portion and 5'-dimethoxytrityl-2'-O-  
methyl-3'-O-phosphoramidite for 5' and 3' wings. The  
standard synthesis cycle is modified by increasing the wait  
5 step after the delivery of tetrazole and base to 600 s  
repeated four times for RNA and twice for 2'-O-methyl. The  
fully protected oligonucleotide is cleaved from the support  
and the phosphate group is deprotected in 3:1 ammonia/ethanol  
at room temperature overnight then lyophilized to dryness.  
10 Treatment in methanolic ammonia for 24 hrs at room  
temperature is then done to deprotect all bases and sample  
was again lyophilized to dryness. The pellet is resuspended  
in 1M TBAF in THF for 24 hrs at room temperature to deprotect  
the 2' positions. The reaction is then quenched with 1M TEAA  
15 and the sample is then reduced to 1/2 volume by rotovac  
before being desalted on a G25 size exclusion column. The  
oligo recovered is then analyzed spectrophotometrically for  
yield and for purity by capillary electrophoresis and by mass  
spectrometry.

20

**[2'-O-(2-Methoxyethyl)]--[2'-deoxy]--[2'-O-(Methoxyethyl)] Chimeric Phosphorothioate Oligonucleotides**

25 [2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(methoxyethyl)] chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.

30

**[2'-O-(2-Methoxyethyl)Phosphodiester]--[2'-deoxy Phosphorothioate]--[2'-O-(2-Methoxyethyl)Phosphodiester] Chimeric Oligonucleotides**

35 [2'-O-(2-methoxyethyl phosphodiester)]--[2'-deoxy phosphorothioate]--[2'-O-(methoxyethyl) phosphodiester] chimeric oligonucleotides are prepared as per the above procedure for the 2'-O-methyl chimeric oligonucleotide with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-

methy1 amidites, oxidization with iodine to generate the phosphodiester internucleotide linkages within the wing portions of the chimeric structures and sulfurization utilizing 3,H-1,2 benzodithiole-3-one 1,1 dioxide (Beaucage Reagent) to generate the phosphorothioate internucleotide linkages for the center gap.

Other chimeric oligonucleotides, chimeric oligonucleosides and mixed chimeric oligonucleotides/oligonucleosides are synthesized according to United States patent 5,623,065, herein incorporated by reference.

#### Example 6

##### Oligonucleotide Isolation

After cleavage from the controlled pore glass column (Applied Biosystems) and deblocking in concentrated ammonium hydroxide at 55°C for 18 hours, the oligonucleotides or oligonucleosides are purified by precipitation twice out of 0.5 M NaCl with 2.5 volumes ethanol. Synthesized oligonucleotides were analyzed by polyacrylamide gel electrophoresis on denaturing gels and judged to be at least 85% full length material. The relative amounts of phosphorothioate and phosphodiester linkages obtained in synthesis were periodically checked by <sup>31</sup>P nuclear magnetic resonance spectroscopy, and for some studies oligonucleotides were purified by HPLC, as described by Chiang et al., *J. Biol. Chem.* **1991**, 266, 18162-18171. Results obtained with HPLC-purified material were similar to those obtained with non-HPLC purified material.

#### Example 7

##### Oligonucleotide Synthesis - 96 Well Plate Format

Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry on an automated synthesizer capable of assembling 96 sequences simultaneously in a standard 96 well format. Phosphodiester internucleotide linkages were afforded by oxidation with aqueous iodine. Phosphorothioate

internucleotide linkages were generated by sulfurization utilizing 3,4,5-benzodithiole-3-one 1,1-dioxide (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial vendors (e.g. PE-Applied Biosystems, Foster City, CA, or Pharmacia, Piscataway, NJ). Non-standard nucleosides are synthesized as per known literature or patented methods. They are utilized as base protected beta-cyanoethyl-diisopropyl phosphoramidites.

Oligonucleotides were cleaved from support and deprotected with concentrated  $\text{NH}_4\text{OH}$  at elevated temperature (55-60°C) for 12-16 hours and the released product then dried in vacuo. The dried product was then re-suspended in sterile water to afford a master plate from which all analytical and test plate samples are then diluted utilizing robotic pipettors.

#### Example 8

##### Oligonucleotide Analysis - 96 Well Plate Format

The concentration of oligonucleotide in each well was assessed by dilution of samples and UV absorption spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96 well format (Beckman P/ACE™ MDQ) or, for individually prepared samples, on a commercial CE apparatus (e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone composition was confirmed by mass analysis of the compounds utilizing electrospray-mass spectroscopy. All assay test plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the compounds on the plate were at least 85% full length.

#### Example 9

##### Cell culture and oligonucleotide treatment

The effect of antisense compounds on target nucleic acid



expression can be tested in any of a variety of cell types provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. The following 5 cell types are provided for illustrative purposes, but other cell types can be routinely used, provided that the target is expressed in the cell type chosen. This can be readily determined by methods routine in the art, for example Northern blot analysis, Ribonuclease protection assays, or RT-PCR.

T-24 cells:

The human transitional cell bladder carcinoma cell line T-24 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). T-24 cells were routinely cultured in complete McCoy's 5A basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analysis, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

A549 cells:

The human lung carcinoma cell line A549 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). A549 cells were routinely cultured in DMEM basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg,

MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence.

NHDF cells:

5 Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville MD). NHDFs were routinely maintained in Fibroblast Growth Medium (Clonetics Corporation, Walkersville MD) supplemented as recommended by the supplier. Cells were maintained for up to 10 passages as  
10 recommended by the supplier.

HEK cells:

Human embryonic keratinocytes (HEK) were obtained from the Clonetics Corporation (Walkersville MD). HEKs were  
15 routinely maintained in Keratinocyte Growth Medium (Clonetics Corporation, Walkersville MD) formulated as recommended by the supplier. Cells were routinely maintained for up to 10 passages as recommended by the supplier.

20 b.END cells:

The mouse brain endothelial cell line b.END was obtained from Dr. Werner Risau at the Max Plank Institute (Bad  
Nauheim, Germany). b.END cells were routinely cultured in  
25 DMEM, high glucose (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-  
30 Primaria #3872) at a density of 3000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium  
35 and oligonucleotide.

Treatment with antisense compounds:

When cells reached 80% confluency, they were treated

with oligonucleotide. For cells grown in 96-well plates, wells were washed once with 200  $\mu$ L OPTI-MEM™-1 reduced-serum medium (Gibco BRL) and then treated with 130  $\mu$ L of OPTI-MEM™-1 containing 3.75  $\mu$ g/mL LIPOFECTIN™ (Gibco BRL) and the  
5 desired concentration of oligonucleotide. After 4-7 hours of treatment, the medium was replaced with fresh medium. Cells were harvested 16-24 hours after oligonucleotide treatment.

The concentration of oligonucleotide used varies from cell line to cell line. To determine the optimal  
10 oligonucleotide concentration for a particular cell line, the cells are treated with a positive control oligonucleotide at a range of concentrations. For human cells the positive control oligonucleotide is ISIS 13920, **TCCGTCATCGCTCCTCAGGG**,  
SEQ ID NO: 1, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls  
15 shown in bold) with a phosphorothioate backbone which is targeted to human H-ras. For mouse or rat cells the positive control oligonucleotide is ISIS 15770, **ATGCATTCTGCCCAAGGA**,  
SEQ ID NO: 2, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls  
20 shown in bold) with a phosphorothioate backbone which is targeted to both mouse and rat c-raf. The concentration of positive control oligonucleotide that results in 80% inhibition of c-Ha-ras (for ISIS 13920) or c-raf (for ISIS 15770) mRNA is then utilized as the screening concentration for new oligonucleotides in subsequent experiments for that  
25 cell line. If 80% inhibition is not achieved, the lowest concentration of positive control oligonucleotide that results in 60% inhibition of H-ras or c-raf mRNA is then utilized as the oligonucleotide screening concentration in subsequent experiments for that cell line. If 60% inhibition  
30 is not achieved, that particular cell line is deemed as unsuitable for oligonucleotide transfection experiments.

#### Example 10

#### Analysis of oligonucleotide inhibition of caspase 7

#### 35 expression

Antisense modulation of caspase 7 expression can be

assayed in a variety of ways known in the art. For example, caspase 7 mRNA levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). Real-time quantitative PCR is presently preferred. RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Northern blot analysis is routine in the art and is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.2.1-4.2.9, John Wiley & Sons, Inc., 1996. Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence Detection System, available from PE-Applied Biosystems, Foster City, CA and used according to manufacturer's instructions.

Protein levels of caspase 7 can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), ELISA or fluorescence-activated cell sorting (FACS). Antibodies directed to caspase 7 can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, MI), or can be prepared via conventional antibody generation methods. Methods for preparation of polyclonal antisera are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.12.1-11.12.9, John Wiley & Sons, Inc., 1997. Preparation of monoclonal antibodies is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.4.1-11.11.5, John Wiley & Sons, Inc., 1997.

Immunoprecipitation methods are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.16.1-10.16.11, John Wiley & Sons, Inc., 1998. Western blot

(immunoblot) analysis is standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.8.1-10.8.21, John Wiley & Sons, Inc., 1997. Enzyme-linked immunosorbent assays (ELISA) are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.2.1-11.2.22, John Wiley & Sons, Inc., 1991.

#### Example 11

##### 10 Poly(A)+ mRNA isolation

Poly(A)+ mRNA was isolated according to Miura et al., *Clin. Chem.*, 1996, 42, 1758-1764. Other methods for poly(A)+ mRNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200  $\mu$ L cold PBS. 60  $\mu$ L lysis buffer (10 mM Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM vanadyl-ribonucleoside complex) was added to each well, the plate was gently agitated and then incubated at room temperature for five minutes. 55  $\mu$ L of lysate was transferred to Oligo d(T) coated 96-well plates (AGCT Inc., Irvine CA). Plates were incubated for 60 minutes at room temperature, washed 3 times with 200  $\mu$ L of wash buffer (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the final wash, the plate was blotted on paper towels to remove excess wash buffer and then air-dried for 5 minutes. 60  $\mu$ L of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70°C was added to each well, the plate was incubated on a 90°C hot plate for 5 minutes, and the eluate was then transferred to a fresh 96-well plate.

Cells grown on 100 mm or other standard plates may be treated similarly, using appropriate volumes of all solutions.

**Example 12****Total RNA Isolation**

Total RNA was isolated using an RNEASY 96™ kit and buffers purchased from Qiagen Inc. (Valencia CA) following the manufacturer's recommended procedures. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200  $\mu$ L cold PBS. 100  $\mu$ L Buffer RLT was added to each well and the plate vigorously agitated for 20 seconds. 100  $\mu$ L of 70% ethanol was then added to each well and the contents mixed by pipetting three times up and down. The samples were then transferred to the RNEASY 96™ well plate attached to a QIAVAC™ manifold fitted with a waste collection tray and attached to a vacuum source. Vacuum was applied for 15 seconds. 1 mL of Buffer RW1 was added to each well of the RNEASY 96™ plate and the vacuum again applied for 15 seconds. 1 mL of Buffer RPE was then added to each well of the RNEASY 96™ plate and the vacuum applied for a period of 15 seconds. The Buffer RPE wash was then repeated and the vacuum was applied for an additional 10 minutes. The plate was then removed from the QIAVAC™ manifold and blotted dry on paper towels. The plate was then re-attached to the QIAVAC™ manifold fitted with a collection tube rack containing 1.2 mL collection tubes. RNA was then eluted by pipetting 60  $\mu$ L water into each well, incubating 1 minute, and then applying the vacuum for 30 seconds. The elution step was repeated with an additional 60  $\mu$ L water.

The repetitive pipetting and elution steps may be automated using a QIAGEN Bio-Robot 9604 (Qiagen, Inc., Valencia CA). Essentially, after lysing of the cells on the culture plate, the plate is transferred to the robot deck where the pipetting, DNase treatment and elution steps are carried out.

**Example 13****Real-time Quantitative PCR Analysis of caspase 7 mRNA Levels**

Quantitation of caspase 7 mRNA levels was determined by real-time quantitative PCR using the ABI PRISM™ 7700 Sequence  
5 Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain  
10 reaction (PCR) products in real-time. As opposed to standard PCR, in which amplification products are quantitated after the PCR is completed, products in real-time quantitative PCR are quantitated as they accumulate. This is accomplished by including in the PCR reaction an oligonucleotide probe that anneals specifically between the forward and reverse PCR  
15 primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE, FAM, or VIC, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, obtained from either Operon  
20 Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence  
25 creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a  
30 sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular intervals by laser optics built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a  
35 series of parallel reactions containing serial dilutions of mRNA from untreated control samples generates a standard

curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

Prior to quantitative PCR analysis, primer-probe sets specific to the target gene being measured are evaluated for their ability to be "multiplexed" with a GAPDH amplification reaction. In multiplexing, both the target gene and the internal standard gene GAPDH are amplified concurrently in a single sample. In this analysis, mRNA isolated from untreated cells is serially diluted. Each dilution is amplified in the presence of primer-probe sets specific for GAPDH only, target gene only ("single-plexing"), or both (multiplexing). Following PCR amplification, standard curves of GAPDH and target mRNA signal as a function of dilution are generated from both the single-plexed and multiplexed samples. If both the slope and correlation coefficient of the GAPDH and target signals generated from the multiplexed samples fall within 10% of their corresponding values generated from the single-plexed samples, the primer-probe set specific for that target is deemed multiplexable. Other methods of PCR are also known in the art.

PCR reagents were obtained from PE-Applied Biosystems, Foster City, CA. RT-PCR reactions were carried out by adding 25  $\mu$ L PCR cocktail (1x TAQMAN<sup>TM</sup> buffer A, 5.5 mM MgCl<sub>2</sub>, 300  $\mu$ M each of dATP, dCTP and dGTP, 600  $\mu$ M of dUTP, 100 nM each of forward primer, reverse primer, and probe, 20 Units RNase inhibitor, 1.25 Units AMPLITAQ GOLD<sup>TM</sup>, and 12.5 Units MuLV reverse transcriptase) to 96 well plates containing 25  $\mu$ L total RNA solution. The RT reaction was carried out by incubation for 30 minutes at 48°C. Following a 10 minute incubation at 95°C to activate the AMPLITAQ GOLD<sup>TM</sup>, 40 cycles of a two-step PCR protocol were carried out: 95°C for 15 seconds (denaturation) followed by 60°C for 1.5 minutes (annealing/extension).

Gene target quantities obtained by real time RT-PCR are normalized using either the expression level of GAPDH, a gene



whose expression is constant, or by quantifying total RNA using RiboGreen™ (Molecular Probes, Inc. Eugene, OR). GAPDH expression is quantified by real time RT-PCR, by being run simultaneously with the target, multiplexing, or separately.

- 5 Total RNA is quantified using RiboGreen™ RNA quantification reagent from Molecular Probes. Methods of RNA quantification by RiboGreen™ are taught in Jones, L.J., et al, *Analytical Biochemistry*, **1998**, 265, 368-374.

In this assay, 175 µL of RiboGreen™ working reagent  
10 (RiboGreen™ reagent diluted 1:2865 in 10mM Tris-HCl, 1 mM EDTA, pH 7.5) is pipetted into a 96-well plate containing 25uL purified, cellular RNA. The plate is read in a CytoFluor 4000 (PE Applied Biosystems) with excitation at 480nm and emission at 520nm.

- 15 Probes and primers to human caspase 7 were designed to hybridize to a human caspase 7 sequence, using published sequence information (GenBank accession number U37448, incorporated herein as SEQ ID NO:3). For human caspase 7 the PCR primers were:

- 20 forward primer: ATTGGTGGTAACAGTGGTAGAGTCAT (SEQ ID NO: 4)  
reverse primer: CCCTTGCTGTGTTTTGTCA (SEQ ID NO: 5) and the PCR probe was: FAM-TTGCACTTGGCAAAAAGAAATCCCAATG-TAMRA (SEQ ID NO: 6) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied  
25 Biosystems, Foster City, CA) is the quencher dye. For human GAPDH the PCR primers were:  
forward primer: CAACGGATTGGTCGTATTGG (SEQ ID NO: 7)  
reverse primer: GGCAACAATATCCACTTTACCAGAGT (SEQ ID NO: 8) and the PCR probe was: 5' JOE-CGCCTGGTCACCAGGGCTGCT- TAMRA 3'  
30 (SEQ ID NO: 9) where JOE (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

- Probes and primers to mouse caspase 7 were designed to hybridize to a mouse caspase 7 sequence, using published  
35 sequence information (GenBank accession number U67321, incorporated herein as SEQ ID NO:10). For mouse caspase 7

the PCR primers were:

forward primer: TGAGGAGGACCACAGCAACTC (SEQ ID NO:11)

reverse primer: TTCCCGTAAATCAGGTCTCTTC (SEQ ID NO: 12) and

the PCR probe was: FAM-CTGCGTCCTGCTGAGCCACGG-TAMRA

- 5 (SEQ ID NO: 13) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

For mouse GAPDH the PCR primers were:

forward primer: GGCAAATTCAACGGCACAGT (SEQ ID NO: 14)

- 10 reverse primer: GGGTCTCGCTCCTGGAGCT (SEQ ID NO: 15) and the PCR probe was: 5' JOE-AAGGCCGAGAATGGGAAGCTTGTCATC- TAMRA 3'

(SEQ ID NO: 16) where JOE (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

15

#### **Example 14**

##### **Northern blot analysis of caspase 7 mRNA levels**

- Eighteen hours after antisense treatment, cell
- 20 monolayers were washed twice with cold PBS and lysed in 1 mL RNazol™ (TEL-TEST "B" Inc., Friendswood, TX). Total RNA was prepared following manufacturer's recommended protocols.
- Twenty micrograms of total RNA was fractionated by electrophoresis through 1.2% agarose gels containing 1.1%
- 25 formaldehyde using a MOPS buffer system (AMRESCO, Inc. Solon, OH). RNA was transferred from the gel to HYBOND™-N+ nylon membranes (Amersham Pharmacia Biotech, Piscataway, NJ) by overnight capillary transfer using a Northern/Southern Transfer buffer system (TEL-TEST "B" Inc., Friendswood, TX).
- 30 RNA transfer was confirmed by UV visualization. Membranes were fixed by UV cross-linking using a STRATALINKER™ UV Crosslinker 2400 (Stratagene, Inc, La Jolla, CA) and then probed using QUICKHYB™ hybridization solution (Stratagene, La Jolla, CA) using manufacturer's recommendations for stringent
- 35 conditions.

To detect human caspase 7, a human caspase 7 specific

probe was prepared by PCR using the forward primer  
ATTGGTGGTAACAGTGGTAGAGTCAT (SEQ ID NO: 4) and the reverse  
primer CCTTGGCTGTGTTTGTCA (SEQ ID NO: 5). To normalize for  
variations in loading and transfer efficiency membranes were  
stripped and probed for human glyceraldehyde-3-phosphate  
dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, CA).

To detect mouse caspase 7, a mouse caspase 7 specific  
probe was prepared by PCR using the forward primer  
TGAGGAGGACCACAGCAACTC (SEQ ID NO:11) and the reverse primer  
TTCCCGTAAATCAGGTCTCTTC (SEQ ID NO: 12). To normalize for  
variations in loading and transfer efficiency membranes were  
stripped and probed for mouse glyceraldehyde-3-phosphate  
dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, CA).

Hybridized membranes were visualized and quantitated  
using a PHOSPHORIMAGER™ and IMAGEQUANT™ Software V3.3  
(Molecular Dynamics, Sunnyvale, CA). Data was normalized to  
GAPDH levels in untreated controls.

## Example 15

### **Antisense inhibition of human caspase 7 expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap**

In accordance with the present invention, a series of  
oligonucleotides were designed to target different regions of  
the human caspase 7 RNA, using published sequences (GenBank  
accession number U37448, incorporated herein as SEQ ID NO: 3,  
GenBank accession number U67206, incorporated herein as SEQ  
ID NO: 17, GenBank accession number U67319, incorporated  
herein as SEQ ID NO: 18, and GenBank accession number U37449,  
incorporated herein as SEQ ID NO: 19). The oligonucleotides  
are shown in Table 1. "Target site" indicates the first (5'-  
most) nucleotide number on the particular target sequence to  
which the oligonucleotide binds. All compounds in Table 1  
are chimeric oligonucleotides ("gapmers") 20 nucleotides in  
length, composed of a central "gap" region consisting of ten

2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human caspase 7 mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present, "N.D." indicates "no data".

Table 1

Inhibition of human caspase 7 mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	%INHIB	SEQ ID NO
135138	5'UTR	3	5	cgqctgggactggcacagtc	45	20
135140	Coding	3	48	aatacagccctgacatctg	50	21
135141	Coding	3	84	atcttcatttgctgaatcct	33	22
135142	Coding	3	94	catccactgaatcttcattt	46	23
135143	Coding	3	104	tctggcttagcatccaactga	90	24
135144	Coding	3	111	ggaccggctctggcttagcat	41	25
135145	Coding	3	138	cttaactgaagaggacggta	53	26
135146	Coding	3	145	tcttcttcttactgaagagg	38	27
135147	Coding	3	168	gatggatcgcatggtagacat	40	28
135148	Coding	3	230	ttgcccagcttttcaaaatt	69	29
135149	Coding	3	332	cttcgggaagcacttgaagag	67	30
135150	Coding	3	338	cccaggcttcggaagcactt	91	31
135151	Coding	3	344	tcaaaaccaggcttcggaa	83	32
135152	Coding	3	354	gacaatcacgtcaaaaccca	62	33
135153	Coding	3	371	caagagcagtcattatagac	69	34
135154	Coding	3	425	gcatttgatggctcctcttc	67	35
135155	Coding	3	496	gtgtgacaccatctttccca	80	36
135156	Coding	3	567	gaagagtttgggtttctcta	44	37
135157	Coding	3	577	cctgaatgaagaagagtttg	36	38
135158	Coding	3	713	taatagccttggaaaccgtga	70	39
135159	Coding	3	716	gagtaatagacctggaacctg	48	40
135160	Coding	3	745	accaggagcctcttccctggg	22	41
135161	Coding	3	751	gcacaaaccaggagcctctt	90	42
135162	Coding	3	778	gctcctccaggatggagcaq	20	43
135163	Coding	3	792	caggtcttttccgtgctcct	80	44
135164	Coding	3	807	gatctgatgatttccaggt	61	45
135165	Coding	3	862	ggctcatcagactgagactca	63	46
135166	Coding	3	911	gtgagcatggagaccacaca	82	47
135167	Coding	3	930	actgaagttagagttccttgg	58	48
135168	Stop	3	944	tgatatggctattgactgaa	77	49

	Codon					
135169	3'UTR	3	971	cattgctcttcagctagaat	77	50
135170	3'UTR	3	977	gtgacccattgctcttcagc	71	51
135171	3'UTR	3	1075	agttttcttccctgttgaatc	69	52
135172	3'UTR	3	1116	aaagctctctgaaaattcaga	21	53
135173	3'UTR	3	1229	cagaagtcacttctgttgca	68	54
135174	3'UTR	3	1237	gcttttctccagaagtcactt	62	55
135175	3'UTR	3	1265	ccaccaattgcaagtgacac	79	56
135176	3'UTR	3	1268	ttaccaccaattgcaagtgga	72	57
135177	3'UTR	3	1363	acattctgcaataaagagca	73	58
135178	3'UTR	3	1370	aataccacattctgcaata	60	59
135179	3'UTR	3	1372	tcaataccacattctgcaa	79	60
135180	3'UTR	3	1407	tgccctaaagccaatgaaaaa	42	61
135181	3'UTR	3	1452	agctttttctcctctaactc	77	62
135182	3'UTR	3	1504	atgggtttctgttttccagac	87	63
135183	3'UTR	3	1551	tgtaagtcatttccctgtat	75	64
135184	3'UTR	3	1615	attatcctcatagccctggg	71	65
135185	3'UTR	3	1663	tcctctggatgatcccaata	90	66
135186	3'UTR	3	1721	ttccacaatccattgggtgg	66	67
135187	3'UTR	3	1747	ggctttgttcttctcatggc	56	68
135188	3'UTR	3	1781	cttctctgtctcagctcact	39	69
135189	3'UTR	3	1783	ctcttatctgtctcagctca	75	70
135190	3'UTR	3	1803	gataggtgagacaaggtatt	14	71
135191	3'UTR	3	1861	tgaggtggcttccagatagg	62	72
135192	3'UTR	3	1899	tcctgcctcactgtcccttg	65	73
135193	3'UTR	3	1939	tcacattatggggccagggtt	67	74
135194	3'UTR	3	1948	tacttatgttcacattatgg	71	75
135195	3'UTR	3	2006	ctatctgacaccccaatgaa	64	76
135196	3'UTR	3	2069	atttttaagctataacagat	25	77
135197	3'UTR	3	2077	ccagatataatttttaagcta	86	78
135198	3'UTR	3	2109	tcaaaataaggcttgggaata	56	79
135199	3'UTR	3	2290	aagcaaccacatttttaatac	53	80
135200	5'UTR	17	78	cgcacgcatagtccctggag	42	81
135201	5'UTR	17	135	ctggctcgggatagtcgcgt	35	82
135202	5'UTR	17	144	gcaccgcccttgctcggga	0	83
135203	5'UTR	17	220	cctttagcgaaaagcgaaact	29	84
135204	5'UTR	17	319	gtctttttccaaagtgcact	42	85
135205	5'UTR	17	347	aaatcccactgtaccacgcg	32	86
135206	5'UTR	18	41	atgtatcccttccctccgagg	1	87
135207	5'UTR	18	123	atacttttctcctccctgtcc	0	88
135208	5'UTR	18	126	cttatacttttctcctccctg	0	89
135209	Coding	18	597	gcgcctcggcatctttgtct	62	90
135210	5'UTR	19	14	tgaagggaagtaccaccacag	25	91
135211	Start Codon	19	130	catcccaacaggctgatccg	41	92
135212	Start Codon	19	135	tctgccatccacaaggctg	49	93
135213	Coding	19	574	catatttttcttctccatggc	85	94
135214	Coding	19	634	agggttgagctcttccgagat	36	95
135215	Stop Codon	3	955	gaatgtaccctgatatggc	65	96

As shown in Table 1, SEQ ID NOs 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40,

42, 44, 45, 46, 47, 48, 49, 50, 51, 52, 54, 55, 56, 57, 58,  
59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 72, 73, 74,  
75, 76, 78, 79, 80, 81, 82, 85, 86, 90, 92, 93, 94, 95 and 96  
demonstrated at least 30% inhibition of human caspase 7  
expression in this assay and are therefore preferred. The  
target sites to which these preferred sequences are  
complementary are herein referred to as "active sites" and  
are therefore preferred sites for targeting by compounds of  
the present invention.

10

#### Example 16

**Antisense inhibition of mouse caspase 7 expression by  
chimeric phosphorothioate oligonucleotides having 2'-MOE  
wings and a deoxy gap.**

15

In accordance with the present invention, a second  
series of oligonucleotides were designed to target different  
regions of the mouse caspase 7 RNA, using published sequences  
(GenBank accession number U67321, incorporated herein as SEQ  
ID NO: 10, GenBank accession number Y13088, incorporated  
herein as SEQ ID NO: 97, and GenBank accession number  
AI116328, incorporated herein as SEQ ID NO: 98). The  
oligonucleotides are shown in Table 2. "Target site"  
indicates the first (5'-most) nucleotide number on the  
particular target sequence to which the oligonucleotide  
binds. All compounds in Table 2 are chimeric  
oligonucleotides ("gapmers") 20 nucleotides in length,  
composed of a central "gap" region consisting of ten 2'-  
deoxynucleotides, which is flanked on both sides (5' and 3'  
directions) by five-nucleotide "wings". The wings are  
composed of 2'-methoxyethyl (2'-MOE)nucleotides. The  
internucleoside (backbone) linkages are phosphorothioate  
(P=S) throughout the oligonucleotide. All cytidine residues  
are 5-methylcytidines. The compounds were analyzed for their  
effect on mouse caspase 7 mRNA levels by quantitative real-  
time PCR as described in other examples herein. Data are  
averages from two experiments. If present, "N.D." indicates

35

SEQUENCE LISTING

"no data".

Table 2

- 5 Inhibition of mouse caspase 7 mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	%INHIB	SEQ ID NO
135156	Coding	10	1108	gaagagtttgggtttctcta	46	37
135157	Coding	10	1118	cetgaatgaagaagagtttg	46	38
135460	5'UTR	97	40	gtgcctgcagcccggaag	33	99
135461	5'UTR	97	95	tgaagcaaaaagtcacgag	66	100
135462	Start Codon	97	182	atcatcgggtcatcggtccca	54	101
135463	Coding	97	413	cgttttgtcgaagttcttgt	74	102
135464	Coding	97	419	acctgtcgttttgtcgaagt	76	103
135465	Coding	97	425	gtccataacctgtcgtttgt	65	104
135466	Coding	97	446	tttgtccgtccattccgga	78	105
135467	Coding	97	474	ggaagcacttgaagaggccc	82	106
135468	Coding	97	598	ctcagcaggacgcagggcga	68	107
135469	Coding	97	603	cgtgggtcagcaggacgcag	80	108
135470	Coding	97	665	atgagctgtcagatccttta	72	109
135471	Coding	97	672	ccctaaaatgagctgtcaga	62	110
135472	Coding	97	679	cggctcctccctaaaatgagc	80	111
135473	Coding	97	686	tttgcatcgggtctccctaa	85	112
135474	Coding	97	692	cagggttttgcatcgggtctc	81	113
135475	Coding	97	772	ggccccgagtcagcctggat	62	114
135476	Coding	97	847	accgtggagtaagcaaaag	53	115
135477	Coding	97	852	ctggaacctggagtaagca	47	116
135478	Coding	97	902	ggcctgcacaaaccaggagc	90	117
135479	Coding	97	909	agcagagggcctgcacaaac	64	118
135480	Coding	97	917	caggatggagcagaggccct	19	119
135481	Coding	97	922	tcattcaggatggagcagag	46	120
135482	Coding	97	929	gccatgctcattcaggatgg	60	121
135483	Coding	97	937	aggctccttgccatgctcatt	64	122
135484	Coding	97	959	ggtcaggatctgcgatgatct	81	123
135485	Coding	97	967	ttcaccctggtcaggatctg	62	124
135486	Coding	97	972	tgtcgttccacctggtcagg	78	125
135487	Coding	97	982	ctggccacctgtcgtttcac	77	126
135488	Coding	97	1003	tcagactgggactcgaagtg	72	127
135489	Coding	97	1057	gtgagcatggacaccataca	62	128
135490	Coding	97	1064	ctctttgtgagcatggaca	70	129
135491	Stop Codon	97	1090	aagggtggctcaacggctgaa	82	130
135492	3'UTR	97	1169	tctggatattttgagagcat	86	131
135493	3'UTR	97	1175	aacattttctggatattttga	71	132
135494	3'UTR	97	1263	cttagagggaagagcatgaag	58	133
135495	3'UTR	97	1330	ggatcatgctcacagcatgca	72	134
135496	3'UTR	97	1446	agcagccacctttgccaatg	71	135
135497	3'UTR	97	1461	tgagaacaaccactcagcag	67	136
135498	3'UTR	97	1480	aaaatctgcacctgaagcact	68	137

135499	3'UTR	97	1539	ttctatatactggaatatta	73	138
135500	3'UTR	97	1599	attccctggtattgacccttc	79	139
135501	3'UTR	97	1606	taagtgaattccctggtattg	76	140
135502	3'UTR	97	1621	tctcaattaccgcgagtgaat	88	141
135503	3'UTR	97	1719	tctagaaggctgacctgcca	87	142
135504	3'UTR	97	1727	ttctagtctctagaaggcttg	61	143
135505	3'UTR	97	1806	tgactcagttcttggtctggt	79	144
135506	3'UTR	97	1850	ggcgtggtggtataggccac	69	145
135507	3'UTR	97	1879	ttggctttcaggcaccttgg	83	146
135508	3'UTR	97	1887	ctcactagtgtggctttcagg	81	147
135509	3'UTR	97	1897	tccctaggccctcactagtt	53	148
135510	3'UTR	97	1917	tgtagcacagggtagcatgg	75	149
135511	3'UTR	97	2004	acgccagaggacatggttat	69	150
135512	3'UTR	97	2033	cagtgaaatagtgaatgaata	54	151
135513	3'UTR	97	2071	actcttcagcacctgtgtgg	90	152
135514	3'UTR	97	2081	gaaatgggttactcttcagg	77	153
135515	3'UTR	97	2096	tcttgtatatacaaatgaaat	51	154
135516	3'UTR	97	2180	cagaggtaacccaaagattat	69	155
135517	3'UTR	97	2211	agatcttccgctaaaattgtc	79	156
135518	3'UTR	97	2304	aaataggaaacatctgtcaac	65	157
135519	5'UTR	10	41	gtatgaaaattctctgtgatt	29	158
135520	5'UTR	10	86	tttccacctgccaattctcac	51	159
135521	5'UTR	10	162	gtgaaaagtgtgctactgttg	40	160
135522	5'UTR	10	203	tgaaaactgcagatttgatg	29	161
135523	5'UTR	10	268	atagccatgtaatatataatgt	31	162
135524	5'UTR	10	311	ttcaatcactgtagtctaaag	27	163
135525	5'UTR	10	324	ctgtacgaagaacttcaatc	37	164
135526	5'UTR	10	331	ttgatggctgtacgaagaac	50	165
135527	5'UTR	10	393	aaggaacaccttttcatgcct	19	166
135528	Start Codon	10	456	attcaaaagcagcagagaggg	0	167
135529	Start Codon	10	471	acttgggctgtgaacattca	24	168
135530	Start Codon	10	474	aacacttgggctgtgaacat	28	169
135531	Coding	10	484	ggactctgagaacacttggg	29	170
135532	Coding	10	508	gggcagcaactcagtttttg	0	171
135533	Exon	98	239	ctgcggtccccaccagtggtg	67	172
135534	Exon	98	313	gacctccacaaaacctcgctt	0	173
135535	Exon	98	320	cattccggacctccacaaac	15	174

- As shown in Table 2, SEQ ID NOs 37, 38, 100, 101, 102,
- 5 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 159, 160, 164, 165 and 172
- 10 demonstrated at least 35% inhibition of mouse caspase 7 expression in this experiment and are therefore preferred.



The target sites to which these preferred sequences are complementary are herein referred to as "active sites" and are therefore preferred sites for targeting by compounds of the present invention.

5

**Example 17****Western blot analysis of caspase 7 protein levels**

Western blot analysis (immunoblot analysis) is carried out using standard methods. Cells are harvested 16-20 h after oligonucleotide treatment, washed once with PBS, suspended in Laemmli buffer (100 ul/well), boiled for 5 minutes and loaded on a 16% SDS-PAGE gel. Gels are run for 1.5 hours at 150 V, and transferred to membrane for western blotting. Appropriate primary antibody directed to caspase 7 is used, with a radiolabelled or fluorescently labeled secondary antibody directed against the primary antibody species. Bands are visualized using a PHOSPHORIMAGER™ (Molecular Dynamics, Sunnyvale CA).

20

**What is claimed is:**

1. A compound 8 to 50 nucleobases in length targeted to a nucleic acid molecule encoding caspase 7, wherein said compound specifically hybridizes with and inhibits the expression of caspase 7.

2. The compound of claim 1 which is an antisense oligonucleotide.

3. The compound of claim 2 wherein the antisense oligonucleotide has a sequence comprising SEQ ID NO: 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 42, 44, 45, 46, 47, 48, 49, 50, 51, 52, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 72, 73, 74, 75, 76, 78, 79, 80, 81, 82, 85, 86, 90, 92, 93, 94, 95, 96, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 159, 160, 164, 165 or 172.

4. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified internucleoside linkage.

5. The compound of claim 4 wherein the modified internucleoside linkage is a phosphorothioate linkage.

6. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified sugar moiety.

7. The compound of claim 6 wherein the modified sugar moiety is a 2'-O-methoxyethyl sugar moiety.

8. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified nucleobase.

9. The compound of claim 8 wherein the modified nucleobase is a 5-methylcytosine.

10. The compound of claim 2 wherein the antisense oligonucleotide is a chimeric oligonucleotide.

11. A compound 8 to 50 nucleobases in length which specifically hybridizes with at least an 8-nucleobase portion of an active site on a nucleic acid molecule encoding caspase 7.

12. A composition comprising the compound of claim 1 and a pharmaceutically acceptable carrier or diluent.

13. The composition of claim 12 further comprising a colloidal dispersion system.

14. The composition of claim 12 wherein the compound is an antisense oligonucleotide.

15. A method of inhibiting the expression of caspase 7 in cells or tissues comprising contacting said cells or tissues with the compound of claim 1 so that expression of caspase 7 is inhibited.

16. A method of treating an animal having a disease or condition associated with caspase 7 comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of claim 1 so that expression of caspase 7 is inhibited.

17. The method of claim 16 wherein the disease or condition is an inflammatory condition.

18. The method of claim 16 wherein the disease or condition is a hyperproliferative disorder.

19. The method of claim 18 wherein the hyperproliferative disorder is cancer.

20. The method of claim 16 wherein the disease or condition is a bone metabolism or cholesterol disorder.

**ABSTRACT**

5

Antisense compounds, compositions and methods are provided for modulating the expression of caspase 7. The compositions comprise antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding caspase 7. Methods of using these compounds for modulation of caspase 7 expression and for treatment of diseases associated with expression of caspase 7 are provided.

10

## COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; and

I verily believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: **Antisense Modulation of caspase 7 Expression** the specification of which:

(XX) is attached hereto.

( ) was filed on \_\_\_\_\_ as Application Serial No. \_\_\_\_\_ and was amended on \_\_\_\_\_ (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information known to be material to the patentability of this application in accordance with 37 CFR § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a-d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of any application on which priority is claimed:

Country	Number	Date Filed	Priority Claimed	
			Yes	No
			Yes	No
			Yes	No

I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information known to be material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

Application Serial No.	Filing Date	Status (pending, patented)

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States provisional application(s) listed below:

Provisional Application No.	Filing Date



DOCKET NO.: RTS-0201

I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: Herb Boswell, Registration No. 27,311; Laurel Spear Bernstein, Registration No. 37,280; Neil S. Bartfeld, Registration No. 39,901; and April C. Logan, Registration No. 33,950, of Isis Pharmaceuticals, Inc.; and Jane Massey Licata, Registration No. 32,257, and Kathleen A. Tyrrell, Registration No. 38,350 of the firm of Law Offices of Jane Massey Licata, 66 East Main Street, Marlton NJ 08053.

Address all telephone calls and correspondence to:

Jane Massey Licata  
Law Offices of Jane Massey Licata  
66 East Main Street  
Marlton NJ 08053  
(856) 810-1515

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

1	Full Name: Hong Zhang	Inventor's Signature: 	Date: 9-6-00
	Residence: 3339 Cadencia St Carlsbad CA 92009	Citizenship: USA	
	Post Office Address:same as above		
2	Full Name: Andrew T. Watt	Inventor's Signature: 	Date: 09/07/00
	Residence: 1500 Shadowridge Dr.. Apt. 154 Vista CA 92083	Citizenship: USA	
	Post Office Address:same as above		

## SEQUENCE LISTING

<110> Hong Zhang  
Andrew T. Watt

<120> ANTISENSE MODULATION OF CASPASE 7 EXPRESSION

<130> RTS-0201

<160> 174

<210> 1

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 1

tccgtcatcg ctctcaggg

20

<210> 2

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 2

atgcattctg cccccaagga

20

<210> 3

<211> 2309

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

&lt;222&gt; (44) ... (955)

&lt;400&gt; 3

gagagactgt gccagtcacca gccgccttac cgccgtggga acg atg gca gat gat	55
Met Ala Asp Asp	
1	
cag ggc tgt att gaa gag cag ggg gtt gag gat tca gca aat gaa gat	103
Gln Gly Cys Ile Glu Glu Gln Gly Val Glu Asp Ser Ala Asn Glu Asp	
5 10 15 20	
tca gtg gat gct aag cca gac cgg tcc tcg ttt gta cgg tcc ctc ttc	151
Ser Val Asp Ala Lys Pro Asp Arg Ser Ser Phe Val Pro Ser Leu Phe	
25 30 35	
agt aag aag aag aaa aat gtc acc atg cga tcc atc aag acc acc cgg	199
Ser Lys Lys Lys Lys Asn Val Thr Met Arg Ser Ile Lys Thr Thr Arg	
40 45 50	
gac cga gtg cct aca tat cag tac aac atg aat ttt gaa aag ctg ggc	247
Asp Arg Val Pro Thr Tyr Gln Tyr Asn Met Asn Phe Glu Lys Leu Gly	
55 60 65	
aaa tgc atc ata ata aac aac aag aac ttt gat aaa gtg aca ggt atg	295
Lys Cys Ile Ile Ile Asn Asn Lys Asn Phe Asp Lys Val Thr Gly Met	
70 75 80	
ggc gtt cga aac gga aca gac aaa gat gcc gag gcg ctc ttc aag tgc	343
Gly Val Arg Asn Gly Thr Asp Lys Asp Ala Glu Ala Leu Phe Lys Cys	
85 90 95 100	
ttc cga agc ctg ggt ttt gac gtg att gtc tat aat gac tgc tct tgt	391
Phe Arg Ser Leu Gly Phe Asp Val Ile Val Tyr Asn Asp Cys Ser Cys	
105 110 115	
gcc aag atg caa gat ctg ctt aaa aaa gct tct gaa gag gac cat aca	439
Ala Lys Met Gln Asp Leu Leu Lys Lys Ala Ser Glu Glu Asp His Thr	
120 125 130	
aat gcc gcc tgc ttc gcc tgc atc ctc tta agc cat gga gaa gaa aat	487
Asn Ala Ala Cys Phe Ala Cys Ile Leu Leu Ser His Gly Glu Glu Asn	
135 140 145	
gta att tat ggg aaa gat ggt gtc aca cca ata aag gat ttg aca gcc	535



Val Ile Tyr Gly Lys Asp Gly Val Thr Pro Ile Lys Asp Leu Thr Ala	
150 155 160	
cac ttt agg ggg gat aga tgc aaa acc ctt tta gag aaa ccc aaa ctc	583
His Phe Arg Gly Asp Arg Cys Lys Thr Leu Leu Glu Lys Pro Lys Leu	
165 170 175 180	
ttc ttc att cag gct tgc cga ggg acc gag ctt gat gat ggc atc cag	631
Phe Phe Ile Gln Ala Cys Arg Gly Thr Glu Leu Asp Asp Gly Ile Gln	
185 190 195	
gcc gac tcg ggg ccc atc aat gac aca gat gct aat cct cga tac aag	679
Ala Asp Ser Gly Pro Ile Asn Asp Thr Asp Ala Asn Pro Arg Tyr Lys	
200 205 210	
atc cca gtg gaa gct gac ttc ctc ttc gcc tat tcc acg gtt cca ggc	727
Ile Pro Val Glu Ala Asp Phe Leu Phe Ala Tyr Ser Thr Val Pro Gly	
215 220 225	
tat tac tcg tgg agg agc cca gga aga ggc tcc tgg ttt gtg caa gcc	775
Tyr Tyr Ser Trp Arg Ser Pro Gly Arg Gly Ser Trp Phe Val Gln Ala	
230 235 240	
ctc tgc tcc atc ctg gag gag cac gga aaa gac ctg gaa atc atg cag	823
Leu Cys Ser Ile Leu Glu Glu His Gly Lys Asp Leu Glu Ile Met Gln	
245 250 255 260	
atc ctc acc agg gtg aat gac aga gtt gcc agg cac ttt gag tct cag	871
Ile Leu Thr Arg Val Asn Asp Arg Val Ala Arg His Phe Glu Ser Gln	
265 270 275	
tct gat gac cca cac ttc cat gag aag aag cag atc ccc tgt gtg gtc	919
Ser Asp Asp Pro His Phe His Glu Lys Lys Gln Ile Pro Cys Val Val	
280 285 290	
tcc atg ctc acc aag gaa ctc tac ttc agt caa tag ccatatcagg	965
Ser Met Leu Thr Lys Glu Leu Tyr Phe Ser Gln	
295 300	
ggtacattct agctgagaag caatgggtca ctcattaatg aatcacattt ttttatgtc	1025
ttgaaatatt cagaaattct ccaggatttt aatttcagga aaatgtattg attcaacagg	1085
gaagaaactt tctggtgctg tcttttgctc tctgaatttt cagagacttt tttataatgt	1145

tatcatttg gtgactgtgt aactttctct taagattaat ttctcttttg tatgtctgtt 1205  
 acctgttaa tagacttaac acatgcaaca gaagtgaact ctggagaaag ctcattgctg 1265  
 tgtccactgc aattggtggt aacagtggtg gagtcattgt tgcacttgcc aaaaagaatc 1325  
 ccaatgtttg acaaaaacaca gccaaagggg tatttactgc tctttattgc agaattgtgg 1385  
 tattgagtgt gatttgaatg atttttcatt ggcttagggc agattttcat gcaaaagtgc 1445  
 tcatatgagt tagaggagaa aaagcttaac gattctgata tgtatccatc aggatccagt 1505  
 ctggaaaaca gaaaccattc taggtgtttc aacagagggg gtttaataca ggaaattgac 1565  
 ttacatagat gataaaagag aagccaaaca gcaagaagct gttaccacac ccagggttat 1625  
 gaggataatg ggaagaggtt tggtttcctg tgtccagtag tgggatcacc cagaggagct 1685  
 ggaaccatgg tgggggctgc ctagtgggag ttaggaccac caatggattg tggaaaatgg 1745  
 agccatgaca agaacaaagc cactgactga gatggagtga gctgagacag ataagagaat 1805  
 acctgtctc acctatcctg ccctcacatc ttccaccagc accttactgc ccaggccat 1865  
 ctggaagcca cctcaccaag gaccttgaa gagcaaggga cagtgaggca ggagaagaac 1925  
 aagaaatgga tgtaagcctg gcccataatg tgaacataag taactactaa tgctcaacaa 1985  
 tttatccatt caatcattta ttcatgggtt tgtcagatag tctatgtatg tgtaaaacaa 2045  
 tctgttttgg ctttatgtgc aaaatctggt atagctttaa aatatatctg gaacttttta 2105  
 gattattcca agccttattt tgagtaaata ttgttactt ttatgtctat aagtgaggaa 2165  
 gagtttatgg caaagatttt tggcactttg tttcaagat ggtgttatct tttgaattct 2225  
 tgataaatga ctgttttttt ctgcctaata gtaactgggt aaaaacaaa tgttcatatt 2285  
 tattgattaa aaatgtggtt gctt 2309

&lt;210&gt; 4

&lt;211&gt; 26

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; PCR Primer

&lt;400&gt; 4

attggtggta acagtggtag agtcoat

26

&lt;210&gt; 5

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; PCR Primer

&lt;400&gt; 5

cccttggtg tggtttgtca

20

&lt;210&gt; 6

&lt;211&gt; 27

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; PCR Probe

&lt;400&gt; 6

ttgcacttg caaaaagaat cccaatg

27

&lt;210&gt; 7

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; PCR Primer

&lt;400&gt; 7

caacggattt ggtcgtattg g

21

&lt;210&gt; 8

&lt;211&gt; 26

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; PCR Primer

&lt;400&gt; 8

ggcaacaata tccactttac cagagt 26

&lt;210&gt; 9

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; PCR Probe

&lt;400&gt; 9

cgcttggtca ccagggtgc t 21

&lt;210&gt; 10

&lt;211&gt; 2006

&lt;212&gt; DNA

&lt;213&gt; Mus musculus

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (474)...(1496)

&lt;400&gt; 10

agctcagtg ggctgatgtg tactgcacat ttaaaaaaaaa aatcacagga attttcatac 60

aatgaataaa accacaacaa tacatgtaga attggcaggt ggaaaagagc cagcaagggc 120

tcaaaactaat cactcacttt ccctcttcag catagttaa ccaacagtag cacactttca 180

cctacaaatc ttaaagtagc tccatcaaat ctgcagtttt cacattatgtg aaaatgtctg 240

tcacataggt acaaatcttg aatcatcaca ttatattaca tggctattct aggtcatcta 300

tagatcagat cttgactac agtgattgaa gttcttcgta cagccatcaa aaaggacac 360

atgatcatta cctactgtta gtcacatct aaaggcatga aaaggtttcc tttttttcaa 420



aaa gcc tct gag gag gac cac agc aac tcg gcc tgc ttc gcc tgc gtc	1004
Lys Ala Ser Glu Glu Asp His Ser Asn Ser Ala Cys Phe Ala Cys Val	
165 170 175	
ctg ctg agc cac ggg gaa gag gac ctg att tac ggg aaa gat ggc gtg	1052
Leu Leu Ser His Gly Glu Glu Asp Leu Ile Tyr Gly Lys Asp Gly Val	
180 185 190	
aca ccc ata aag gat ctg aca gct cat ttt agg gga gac cga tgc aaa	1100
Thr Pro Ile Lys Asp Leu Thr Ala His Phe Arg Gly Asp Arg Cys Lys	
195 200 205	
acc ctg tta gag aaa ccc aaa ctc ttc ttc att cag gca tgc cga ggg	1148
Thr Leu Leu Glu Lys Pro Lys Leu Phe Phe Ile Gln Ala Cys Arg Gly	
210 215 220 225	
acg gag ctc gac gat gga atc cag gct gac tcg ggg ccc atc aac gac	1196
Thr Glu Leu Asp Asp Gly Ile Gln Ala Asp Ser Gly Pro Ile Asn Asp	
230 235 240	
att gac gct aat ccc cgc aac aag atc cgg gtg gaa gcc gac ttc ctc	1244
Ile Asp Ala Asn Pro Arg Asn Lys Ile Pro Val Glu Ala Asp Phe Leu	
245 250 255	
ttt gct tac tcc acg gtt cca ggt tat tac tca tgg agg aac cca ggg	1292
Phe Ala Tyr Ser Thr Val Pro Gly Tyr Tyr Ser Trp Arg Asn Pro Gly	
260 265 270	
aaa ggc tcc tgg ttt gtg cag gcc ctc tgc tcc atc ctg aat gag cat	1340
Lys Gly Ser Trp Phe Val Gln Ala Leu Cys Ser Ile Leu Asn Glu His	
275 280 285	
ggc aag gac ctc gag atc atg cag atc ctg acc agg gtg aac gac agg	1388
Gly Lys Asp Leu Glu Ile Met Gln Ile Leu Thr Arg Val Asn Asp Arg	
290 295 300 305	
gtg gcc agg cac ttc gag tcc cag tct gat gat cca cgc ttc aac gag	1436
Val Ala Arg His Phe Glu Ser Gln Ser Asp Asp Pro Arg Phe Asn Glu	
310 315 320	
aag aag cag atc cgg tgt atg gtg tcc atg ctc acc aaa gag ctg tac	1484
Lys Lys Gln Ile Pro Cys Met Val Ser Met Leu Thr Lys Glu Leu Tyr	
325 330 335	

ttc agc cgt tga ccacccttca gctgagaacc tgccgccgtt cgttgatgaa 1536  
Phe Ser Arg

340

tccagttttt atttttat tttt tgttccgatg ctctcaaaat atccagaaat gttgagggga 1596

tttaatttca ggaaagtcta gatttttttt ttttgtttaa taactttgtt catctgatga 1656

cttcacgtctc ttctctaaag gttgatattcc tgtttctggt tctttttttc tttgtcgtct 1716

cgctgagtg c atgctgtgag catgacctct ggagaagaca ttggcaatga cgtctcagtt 1776

gaacttggca aagagaatcc cagctcttga tgaaagaata cagctgcgac acctgttggc 1836

ctccattggc aaaggtggct gctgagtggt tgttctcagt ggcttagggc agatttttaa 1896

gccgaccttc ccaggtggct gagagaagac gacagttaat attccagtat atagaacca 1956

atccagaaaa taagccatcc taggaatata ggtgcagaag ggtcaataca 2006

<210> 11

<211> 21

<212> DNA

<213> Artificial Sequence

<223> PCR Primer

<400> 11

tgaggaggac cacagcaact c 21

<210> 12

<211> 23

<212> DNA

<213> Artificial Sequence

<223> PCR Primer

<400> 12

ttcccgtaaa tcaggtctctc ttc 23

<210> 13

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; PCR Probe

&lt;400&gt; 13

ctgcgtcctg ctgagccacg g

21

&lt;210&gt; 14

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; PCR Primer

&lt;400&gt; 14

ggcaaattca acggcacagt

20

&lt;210&gt; 15

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; PCR Primer

&lt;400&gt; 15

gggtctcgct cctggaagct

20

&lt;210&gt; 16

&lt;211&gt; 27

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; PCR Probe

&lt;400&gt; 16

aaggccgaga atgggaagct tgtcatc

27

&lt;210&gt; 17



&lt;211&gt; 2661

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (387)...(1298)

&lt;400&gt; 17

cgcccgcgcg cgggctcaac ttgtagagc gagggggccaa ctggcgagag cgcgcgccca 60

gctttgcaga gagcgccctc cagggactat gcgtgcgggg acacgggtcg ctttgggctc 120

ttccaccctt gcgagcgca ctatcccgag ccagggcgcg tgcaagcccc gcccgccct 180

accagggcg gctcctcctt ccgcagcgcc gagactttta gtttcgcttt cgctaaaggg 240

gccccagacc cttgctgcgg agcgacggag agagactgtg ccagtcacag cggccctacc 300

gccgtgggaa cggcaggaag tggcacttgg aaaagaacac cagctgcggt ggtagcagtg 360

ggatttgtgc ttcttatgtt acccag atg gca gat gat cag ggc tgt att gaa 413

Met Ala Asp Asp Gln Gly Cys Ile Glu

1

5

gag cag ggg gtt gag gat tca gca aat gaa gat tca gtg gat gct aag 461

Glu Gln Gly Val Glu Asp Ser Ala Asn Glu Asp Ser Val Asp Ala Lys

10

15

20

25

cca gac cgg tcc tcg ttt gta ccg tcc ctc ttc agt aag aag aag aaa 509

Pro Asp Arg Ser Ser Phe Val Pro Ser Leu Phe Ser Lys Lys Lys Lys

30

35

40

aat gtc acc atg cga tcc atc aag acc acc cgg gac cga gtg cct aca 557

Asn Val Thr Met Arg Ser Ile Lys Thr Thr Arg Asp Arg Val Pro Thr

45

50

55

tat cag tac aac atg aat ttt gaa aag ctg ggc aaa tgc atc ata ata 605

Tyr Gln Tyr Asn Met Asn Phe Glu Lys Leu Gly Lys Cys Ile Ile Ile

60

65

70

aac aac aag aac ttt gat aaa gtg aca ggt atg ggc gtt cga aac gga 653

Asn Asn Lys Asn Phe Asp Lys Val Thr Gly Met Gly Val Arg Asn Gly

75

80

85

aca gac aaa gat gcc gag gcg ctc ttc aag tgc ttc cga agc ctg ggt 701  
 Thr Asp Lys Asp Ala Glu Ala Leu Phe Lys Cys Phe Arg Ser Leu Gly  
 90 95 100 105

ttt gac gtg att gtc tat aat gac tgc tct tgt gcc aag atg caa gat 749  
 Phe Asp Val Ile Val Tyr Asn Asp Cys Ser Cys Ala Lys Met Gln Asp  
 110 115 120

ctg ctt aaa aaa gct tct gaa gag gac cat aca aat gcc gcc tgc ttc 797  
 Leu Leu Lys Lys Ala Ser Glu Glu Asp His Thr Asn Ala Ala Cys Phe  
 125 130 135

gcc tgc atc ctc tta agc cat gga gaa gaa aat gta att tat ggg aaa 845  
 Ala Cys Ile Leu Leu Ser His Gly Glu Glu Asn Val Ile Tyr Gly Lys  
 140 145 150

gat ggt gtc aca cca ata aag gat ttg aca gcc cac ttt agg ggg gat 893  
 Asp Gly Val Thr Pro Ile Lys Asp Leu Thr Ala His Phe Arg Gly Asp  
 155 160 165

aga tgc aaa acc ctt tta gag aaa ccc aaa ctc ttc ttc att cag gct 941  
 Arg Cys Lys Thr Leu Leu Glu Lys Pro Lys Leu Phe Phe Ile Gln Ala  
 170 175 180 185

tgc cga ggg acc gag ctt gat gat ggc atc cag gcc gac tgc ggg ccc 989  
 Cys Arg Gly Thr Glu Leu Asp Asp Gly Ile Gln Ala Asp Ser Gly Pro  
 190 195 200

atc aat gac aca gat gct aat cct cga tac aag atc cca gtg gaa gct 1037  
 Ile Asn Asp Thr Asp Ala Asn Pro Arg Tyr Lys Ile Pro Val Glu Ala  
 205 210 215

gac ttc ctc ttc gcc tat tcc acg gtt cca ggc tat tac tca tgg agg 1085  
 Asp Phe Leu Phe Ala Tyr Ser Thr Val Pro Gly Tyr Tyr Ser Trp Arg  
 220 225 230

agc cca gga aga ggc tcc tgg ttt gtg caa gcc ctc tgc tcc atc ctg 1133  
 Ser Pro Gly Arg Gly Ser Trp Phe Val Gln Ala Leu Cys Ser Ile Leu  
 235 240 245

gag gag cac gga aaa gac ctg gaa atc atg cag atc ctc acc agg gtg 1181  
 Glu Glu His Gly Lys Asp Leu Glu Ile Met Gln Ile Leu Thr Arg Val  
 250 255 260 265

aat gac aga gtt gcc agg cac ttt gag tct cag tct gat gac cca cac 1229  
 Asn Asp Arg Val Ala Arg His Phe Glu Ser Gln Ser Asp Asp Pro His  
 270 275 280

ttc cat gag aag aag cag atc ccc tgt gtg gtc tcc atg ctc acc aag 1277  
 Phe His Glu Lys Lys Gln Ile Pro Cys Val Val Ser Met Leu Thr Lys  
 285 290 295

gaa ctc tac ttc agt caa tag ccatatcagg ggtacattct agctgagaag 1328  
 Glu Leu Tyr Phe Ser Gln  
 300

caatgggtca ctcatatg aatcacattt ttttatgctc ttgaaatatt cagaaattct 1388

ccaggatttt aatttcagga aaatgtattg attcaacagg gaagaaactt tctgggtgct 1448

gtcttttggt ctctgaattt tcagagactt tttttataat gttattcatt tggtgactgt 1508

gkaactttct cttaagatta attttctctt tgtatgtctg ttaccttggt aatagactta 1568

atacatgcaa cagaagtgc ttctggagaa agctcatggc tgtgtccact gcaattgggtg 1628

gcaacagtgg cagatgcatg tttgcacttg gcaaaaagaa tcccaatggt tgacaaacac 1688

agccaagggg atatttactg ctctttattg cagaatgtgg gtattgagtg tgatttgaat 1748

gatttttcat tggcttaggg cagattttca tgcaaaagt ctcatatgag ttagaggaga 1808

aaaagcttaa tgattctgat atgtatccat caggatccag tctggaaaac agaaaccatt 1868

ctaggtgttt caacagaggg agtttaatac aggaaattga cttacataga tgatataaga 1928

gaacccaac agcaagaagc tgttaccaca cccaggtcta tgaggataat gggaagaggt 1988

ttggtttcct gtgtccagta gtgggatcat ccagaggagc tggaaccatg gtgggggctg 2048

cctagtggga gtaggacca ccaatggatt gtggaaaatg gagccatgac aagaacaaaa 2108

ccactgactg agatggagtg agctgagaca gataagagaa taccttggtc tcacctatcc 2168

tgccctcaca tcttccacca gcaccttact gcccaggcct atctggaagc cacctacca 2228

aggaccttgg aagagcaagg gacagtgagg caggagaaga acaagaaatg gatgtaagcc 2288

tggcccataa tgtgaacata agtaatcact aatgctcaac aatttatcca ttcaatcatt 2348  
 tattcattgg gttgtcagat agtctatgta tgtgtaaaac aatctgtttt ggctttatgt 2408  
 ccaaaatctg ttatagcttt aaaatatatc tggaaacttt tagattattc caagccttat 2468  
 tttgagtaaa tattttgttac ttttagttct ataagtgagg aagagtttat ggcaaagatt 2528  
 tttggcactt tgttttcaag atgggtgttat cttttgaatt cttgataaat gactgttttt 2588  
 ttctgcctaa tagtaactgg ttaaaaaaca aatgttcata tttattgatt aaaaatgtgg 2648  
 ttgcttaatt cct 2661

&lt;210&gt; 18

&lt;211&gt; 2602

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (229) ... (1239)

&lt;400&gt; 18

gtggggtaaa acagcagaga tcaatgagat cagagcacac cctcggagga agggatacat 60

gacaaatgcc tgaacggaga gagggagtga actgtgcaaa cacacagcca ggagttttcc 120

aaggacaggg aggagaaagt ataaggcctg ctgtaccctc gatgcaaaac atgagaaagc 180

cgactgtgcc agtcccagcc gcctaccgc cgtgggaacg atgctgta atg gac tgt 237  
 Met Asp Cys  
 1

gtt ggt tgg cct cca ggc agg aag tgg cac ttg gaa aag aac acc agc 285  
 Val Gly Trp Pro Pro Gly Arg Lys Trp His Leu Glu Lys Asn Thr Ser  
 5 10 15

tgc ggt ggt agc agt ggg att tgt gct tct tat gtt acc cag atg gca 333  
 Cys Gly Gly Ser Ser Gly Ile Cys Ala Ser Tyr Val Thr Gln Met Ala  
 20 25 30 35

gat gat cag ggc tgt att gaa gag cag ggg gtt gag gat tca gca aat 381  
 Asp Asp Gln Gly Cys Ile Glu Glu Gln Gly Val Glu Asp Ser Ala Asn  
                   40                  45                  50

gaa gat tca gtg gat gct aag cca gac cgg tcc tgc ttt gta ccg tcc 429  
 Glu Asp Ser Val Asp Ala Lys Pro Asp Arg Ser Ser Phe Val Pro Ser  
                   55                  60                  65

ctc ttc agt aag aag aag aaa aat gtc acc atg cga tcc atc aag acc 477  
 Leu Phe Ser Lys Lys Lys Lys Asn Val Thr Met Arg Ser Ile Lys Thr  
                   70                  75                  80

acc cgg gac cga gtg cct aca tat cag tac aac atg aat ttt gaa aag 525  
 Thr Arg Asp Arg Val Pro Thr Tyr Gln Tyr Asn Met Asn Phe Glu Lys  
                   85                  90                  95

ctg ggc aaa tgc atc ata ata aac aac aag aac ttt gat aaa gtg aca 573  
 Leu Gly Lys Cys Ile Ile Ile Asn Asn Lys Asn Phe Asp Lys Val Thr  
                   100                  105                  110                  115

ggg atg ggc gtt cga aac gga aca gac aaa gat gcc gag gcg ctc ttc 621  
 Gly Met Gly Val Arg Asn Gly Thr Asp Lys Asp Ala Glu Ala Leu Phe  
                   120                  125                  130

aag tgc ttc cga agc ctg ggt ttt gac gtg att gtc tat aat gac tgc 669  
 Lys Cys Phe Arg Ser Leu Gly Phe Asp Val Ile Val Tyr Asn Asp Cys  
                   135                  140                  145

tct tgt gcc aag atg caa gat ctg ctt aaa aaa gct tct gaa gag gac 717  
 Ser Cys Ala Lys Met Gln Asp Leu Leu Lys Lys Ala Ser Glu Glu Asp  
                   150                  155                  160

cat aca aat gcc gcc tgc ttc gcc tgc atc ctc tta agc cat gga gaa 765  
 His Thr Asn Ala Ala Cys Phe Ala Cys Ile Leu Leu Ser His Gly Glu  
                   165                  170                  175

gaa aat gta att tat ggg aaa gat ggt gtc aca cca ata aag gat ttg 813  
 Glu Asn Val Ile Tyr Gly Lys Asp Gly Val Thr Pro Ile Lys Asp Leu  
                   180                  185                  190                  195

aca gcc cac ttt agg ggg gat aga tgc aaa acc ctt tta gag aaa ccc 861  
 Thr Ala His Phe Arg Gly Asp Arg Cys Lys Thr Leu Leu Glu Lys Pro  
                   200                  205                  210

aaa ctc ttc ttc att cag gct tgc cga ggg acc gag ctt gat gat ggc 909  
 Lys Leu Phe Phe Ile Gln Ala Cys Arg Gly Thr Glu Leu Asp Asp Gly  
 215 220 225

atc cag gcc gac tcg ggg ccc atc aat gac aca gat gct aat cct cga 957  
 Ile Gln Ala Asp Ser Gly Pro Ile Asn Asp Thr Asp Ala Asn Pro Arg  
 230 235 240

tac aag atc cca gtg gaa gct gac ttc ctc ttc gcc tat tcc acg gtt 1005  
 Tyr Lys Ile Pro Val Glu Ala Asp Phe Leu Phe Ala Tyr Ser Thr Val  
 245 250 255

cca ggc tat tac tca tgg agg agc cca gga aga ggc tcc tgg ttt gtg 1053  
 Pro Gly Tyr Tyr Ser Trp Arg Ser Pro Gly Arg Gly Ser Trp Phe Val  
 260 265 270 275

caa gcc ctc tgc tcc atc ctg gag gag cac gga aaa gac ctg gaa atc 1101  
 Gln Ala Leu Cys Ser Ile Leu Glu Glu His Gly Lys Asp Leu Glu Ile  
 280 285 290

atg cag atc ctc acc agg gtg aat gac aga gtt gcc agg cac ttt gag 1149  
 Met Gln Ile Leu Thr Arg Val Asn Asp Arg Val Ala Arg His Phe Glu  
 295 300 305

tct cag tct gat gac cca cac ttc cat gag aag aag cag atc ccc tgt 1197  
 Ser Gln Ser Asp Asp Pro His Phe His Glu Lys Lys Gln Ile Pro Cys  
 310 315 320

gtg gtc tcc atg ctc acc aag gaa ctc tac ttc agt caa tag ccatatcagg 1249  
 Val Val Ser Met Leu Thr Lys Glu Leu Tyr Phe Ser Gln  
 325 330 335

ggtacattct agctgagaag caatgggtca ctcattaatg aatcacattt ttttatgctc 1309

ttgaaatatt cagaaattct ccaggatttt aatttcaggga aaatgtattg attcaacagg 1369

gaagaaactt tctgggtgct gtctttttgtt ctctgaattt tcagagactt tttttataat 1429

gttattcatt tgggtactgt gkaactttct cttaagatta attttctctt tgtatgtctg 1489

ttacctgtgt aatagactta atacatgcaa cagaagtgcac ttctggagaa agctcatggc 1549

tgtgtccact gcaattgggt gcaacagtgg cagagtcatg ttgacacttg gcaaaaagaa 1609

tcccaatggt tgacaaacac agccaagggg atatttactg ctctttattg cagaatgtgg 1669  
 gtattgagtg tgatttgaat gatttttcat tggcttaggg cagattttca tgcaaaagtt 1729  
 ctcatatgag ttagaggaga aaaagcttaa tgattctgat atgtatccat caggatccag 1789  
 tctggaaaac agaaaccatt ctagggtgtt caacagaggg agttaatac aggaaattga 1849  
 cttacataga tgatataaga gaacccaaac agcaagaagc tgttaccaca ccagggtcta 1909  
 tgaggataat gggaagaggt ttgggttctt gtgtccagta gtgggatcat ccaggaggac 1969  
 tggaaccatg gtggggggctg cctagtggga gttaggacca ccaatggatt gtggaaaatg 2029  
 gagccatgac agaacaacaa ccactgactg agatggagtg agctgagaca gataagagaa 2089  
 taccttggtc tcacctatcc tgcctcaca tcttccacca gcaccttact gccaggcct 2149  
 atctggaagc cacctcacca aggaccttg aagagcaagg gacagtgagg caggagaaga 2209  
 acaagaaatg gatgtaagcc tggcccataa tgtgaacata agtaatcact aatgctcaac 2269  
 aatttatcca ttcaatcatt tattcattgg gttgtcagat agtctatgta tgtgtaaaac 2329  
 aatctgtttt ggctttatgt ccaaaatctg ttatagcttt aaaatatatc tggaactttt 2389  
 tagattatco caagccttat ttgagtaaa tatttgttac ttttagttct ataagtgagg 2449  
 aagagtttat ggcaagagatt ttgggacttt tgttttcaag atgggtgttat cttttgaatt 2509  
 cttgataaaa gactgttttt ttctgcctaa tagtaactgg ttaaaaaaca aatgttcata 2569  
 tttattgatt aaaaatgtgg ttgcttaatt cct 2602

&lt;210&gt; 19

&lt;211&gt; 2378

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (147) ... (908)

&lt;400&gt; 19

gcaagctggg ctgctgggtg ggtacttctt tcaaagctga gggagcgtcc tacgcccacg 60

cgcgcaggag ggcgcccccc gcaaagcaac gtctaggaga ccacagtgga tgccacagcg 120

ggcccgaagc ggatcagcct tgtggg atg gca gat gat cag ggc tgt att gaa 173

Met Ala Asp Asp Gln Gly Cys Ile Glu

1

5

gag cag ggg gtt gag gat tca gca aat gaa gat tca gtg gat gct aag 221

Glu Gln Gly Val Glu Asp Ser Ala Asn Glu Asp Ser Val Asp Ala Lys

10

15

20

25

cca gac cgg tcc tcg ttt gta ccg tcc ctc ttc agt aag aag aag aaa 269

Pro Asp Arg Ser Ser Phe Val Pro Ser Leu Phe Ser Lys Lys Lys Lys

30

35

40

aat gtc acc atg cga tcc atc aag acc acc cgg gac cga gtg cct aca 317

Asn Val Thr Met Arg Ser Ile Lys Thr Thr Arg Asp Arg Val Pro Thr

45

50

55

tat cag tac aac atg aat ttt gaa aag ctg ggc aaa tgc atc ata ata 365

Tyr Gln Tyr Asn Met Asn Phe Glu Lys Leu Gly Lys Cys Ile Ile Ile

60

65

70

aac aac aag aac ttt gat aaa gtg aca ggt atg ggc gtt cga aac gga 413

Asn Asn Lys Asn Phe Asp Lys Val Thr Gly Met Gly Val Arg Asn Gly

75

80

85

aca gac aaa gat gcc gag gcg ctc ttc aag tgc ttc cga agc ctg ggt 461

Thr Asp Lys Asp Ala Glu Ala Leu Phe Lys Cys Phe Arg Ser Leu Gly

90

95

100

105

ttt gac gtg att gtc tat aat gac tgc tct tgt gcc aag atg caa gat 509

Phe Asp Val Ile Val Tyr Asn Asp Cys Ser Cys Ala Lys Met Gln Asp

110

115

120

ctg ctt aaa aaa gct tct gaa gag gac cat aca aat gcc gcc tgc ttc 557

Leu Leu Lys Lys Ala Ser Glu Glu Asp His Thr Asn Ala Ala Cys Phe

125

130

135

gcc tgc atc ctc tta agc cat gga gaa gaa aat atg gaa tct tgc tct 605

Ala Cys Ile Leu Leu Ser His Gly Glu Glu Asn Met Glu Ser Cys Ser

140

145

150

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
18



gtc acc cag gct gga gtg cag cgg cgt gat ctc gga aga ctg caa cct 653  
 Val Thr Gln Ala Gly Val Gln Arg Asp Leu Gly Arg Leu Gln Pro  
 155 160 165

cca cct ccc agg ctt gcc gag gga ccg agc ttg atg atg gca tcc agg 701  
 Pro Pro Pro Arg Leu Ala Glu Gly Pro Ser Leu Met Met Ala Ser Arg  
 170 175 180 185

ccg act cgg ggc cca tca atg aca cag atg cta atc ctc gat aca aga 749  
 Pro Thr Arg Gly Pro Ser Met Thr Gln Met Leu Ile Leu Asp Thr Arg  
 190 195 200

tcc cag tgg aag ctg act tcc tct tgg cct att cca cgg ttc cag gct 797  
 Ser Gln Trp Lys Leu Thr Ser Ser Ser Pro Ile Pro Arg Phe Gln Ala  
 205 210 215

att act cgt gga gga gcc cag gaa gag gct cct ggt ttg tgc aag ccc 845  
 Ile Thr Arg Gly Gly Ala Gln Glu Glu Ala Pro Gly Leu Cys Lys Pro  
 220 225 230

tct gct cca tcc tgg agg agc acg gaa aag acc tgg aaa tca tgc aga 893  
 Ser Ala Pro Ser Trp Arg Ser Thr Glu Lys Thr Trp Lys Ser Cys Arg  
 235 240 245

tcc tca cca ggg tga atgacagagt tgccaggcac ttgagttctc agtctgatga 948  
 Ser Ser Pro Gly  
 250

cccacacttc catgagaaga agcagatccc ctgtgtggtc tccatgtctca ccaaggaact 1008

ctacttcagt caatagccat atcaggggta cattctagct gagaagcaat gggtcactca 1068

ttaatgaatc acattttttt atgctcttga aatattcaga aattctccag gattttaatt 1128

tcaggaaaat gtattgattc aacaggaag aaactttctg gtgctgtctt ttgttctctg 1188

aattttcaga gactttttta taatgttatt catttggtga ctgtgtaact ttctcttaag 1248

attaattttc tctttgtatg tctgttacct tgtaataaga cttaatacat gcaacagaag 1308

tgacttctgg agaaagctca tggctgtgtc cactgcaatt ggtggtgaaca gtggtagagt 1368

catgtttgca cttggcaaaa agaatcccaa tgtttgacaa aacacagcca aggggatatt 1428

tactgctctt tattgcagaa tgtgggtatt gagtgtgatt tgaatgatt ttcattggct 1488

tagggcagat ttcatgcaa aagtctcat atgagttaga ggagaaaaag cttaatgatt 1548

ctgatatgta tccatcagga tccagtctgg aaaacagaaa ccattctagg tgtttcaaca 1608

gagggagttt aatacagga attgacttac atagatgata aaagagaagc caaacagcaa 1668

gaagctgtta ccacaccag ggctatgagg ataatgggaa gaggtttggt ttcctgtgtc 1728

cagtagtggg atcatccaga ggagctggaa ccaggtggg ggctgcctag tgggagttag 1788

gaccaccaat ggattgtgga aaatggagcc atgacaagaa caaagccact gactgagatg 1848

gagtgaagct agacagataa gagaatacct tgtctcact atctgcctc cacatcttcc 1908

accagcacct tactgcccag gcctatctgg aagccacctc accaaggacc ttggaagagc 1968

aagggacagt gaggcaggag aagaacaaga aatggatgta agcctggccc ataatgtgaa 2028

cataagtaat cactaatgct caacaattta tccattcaat catttattca ttgggtgtgc 2088

agatagtcta tgtatgtgta aaacaatctg ttttggtttt atgtgcaaaa tctgttatag 2148

ctttaaaata tatctggaac tttttagatt attccaagcc ttattttgag taaatatttg 2208

ttacttttag ttctataagt gaggaagagt ttatggcaaa gatttttggc actttgtttt 2268

caagatggtg ttatcttttg aattcttgat aaatgactgt tttttctgc ctaatagtaa 2328

ctggttaaaa aacaaatgtt catatttatt gattaaaaat gtggttgctt 2378

&lt;210&gt; 20

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 20

cggtcgggac tggcacagtc

<210> 21  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 21  
aatacagccc tgatcatctg 20

<210> 22  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 22  
atcttcattt gctgaatcct 20

<210> 23  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 23  
catccactga atcttcattt 20

<210> 24  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 24  
tctggcttag catccactga 20

<210> 25  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 25  
ggaccgggtct ggcttagcat 20

<210> 26  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 26  
cttactgaag agggacggtg 20

<210> 27  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 27  
tcttcttctt actgaagagg 20

<210> 28  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 28  
gatggatcgc atggtgacat 20

<210> 29  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 29  
ttgccagct tttcaaaatt 20

<210> 30  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 30  
cttcggaagc acttgaagag 20

<210> 31  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 31  
cccaggcttc ggaagcactt 20

<210> 32  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 32  
tcaaaaacca ggcttcggaa 20

<210> 33  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 33  
gacaatcacg tcaaaaccca 20

<210> 34  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 34  
caagagcagt cattatagac 20

<210> 35  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 35  
gcatttgat ggctctcttc 20

<210> 36  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 36  
gtgtgacacc atctttccca 20

<210> 37  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 37  
gaagagtttg ggtttctcta 20

<210> 38  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 38  
cctgaatgaa gaagagtttg 20

<210> 39  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 39  
taatagcctg gaaccgtgga 20

<210> 40  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 40  
gagtaatagc ctggaaccgt 20

<210> 41  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 41  
accaggagcc tcttcctggg 20

<210> 42  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 42  
gcacaaacca ggagcctctt 20

<210> 43  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 43  
gctcctccag gatggagcag 20

<210> 44  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 44  
caggtctttt ccgtgctcct 20



<210> 45  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 45  
gatctgcatg atttccaggt 20

<210> 46  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 46  
ggatcatcaga ctgagactca 20

<210> 47  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 47  
gtgagcatgg agaccacaca 20

<210> 48  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 48  
actgaagtag agttccttgg 20

&lt;210&gt; 49

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 49

tgatattggct attgactgaa

20

&lt;210&gt; 50

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 50

cattgcttct cagctagaat

20

&lt;210&gt; 51

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 51

gtgaccatt gcttctcagc

20

&lt;210&gt; 52

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 52

agtttcttcc ctgttgaatc

20

<210> 53  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 53  
aaagtctctg aaaattcaga 20

<210> 54  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 54  
cagaagtcac ttctgttgca 20

<210> 55  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 55  
gcttttcca gaagtcactt 20

<210> 56  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 56  
ccaccaattg cagtggacac 20

<210> 57  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 57  
ttaccaccaa ttgcagtgga 20

<210> 58  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 58  
acattctgca ataaagagca 20

<210> 59  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 59  
aatacccaca ttctgcaata 20

<210> 60  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 60  
tcaataccca cattctgcaa 20

<210> 61  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 61  
tgc cctaagc caatgaaaaa 20

<210> 62  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 62  
agctttttct cctctaactc 20

<210> 63  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 63  
atgggtttctg ttttcagac 20

<210> 64  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 64  
tgtaagtcaa tttcctgtat 20

<210> 65  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 65  
attatcctca tagccctggg 20

<210> 66  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 66  
tcctctggat gatcccacta 20

<210> 67  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 67  
ttccacaatc cattggtggt 20

<210> 68  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 68  
ggctttgttc ttgtcatggc 20

<210> 69  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 69  
cttatctgtc tcagctcact 20

<210> 70  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 70  
ctcttatctg tctcagctca 20

<210> 71  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 71  
gataggtgag acaaggtatt 20

<210> 72  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 72  
tgaggtggct tccagatagg 20

<210> 73  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 73  
tcctgectca ctgtcccttg

20

<210> 74  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 74  
tcacattatg ggccaggctt

20

<210> 75  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 75  
tacttatggtt cacattatgg

20

<210> 76  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 76  
ctatctgaca acccaatgaa

20



&lt;210&gt; 77

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 77

atttttaaagc tataacagat

20

&lt;210&gt; 78

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 78

ccagatatat tttaaagcta

20

&lt;210&gt; 79

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 79

tcaaaaataag gcttgaata

20

&lt;210&gt; 80

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 80

aagcaaccac atttttaatc

20

<210> 81  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 81  
cgcacgcata gtcctggag 20

<210> 82  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 82  
ctggctcggg atagtgcgct 20

<210> 83  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 83  
gcaccgcccc tggctcgga 20

<210> 84  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 84  
cctttagcga aagcgaaact 20

<210> 85  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 85  
gttctttttcc aagtgcact 20

<210> 86  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 86  
aatcccaact gctaccaccg 20

<210> 87  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 87  
atgtatccct tcctccgagg 20

<210> 88  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 88  
atactttctc etccctgtcc 20

<210> 89  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 89  
cttatacttt ctctccctg

20

<210> 90  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 90  
gcgcctcggc atctttgtct

20

<210> 91  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 91  
tgaaggaagt acccaccag

20

<210> 92  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 92  
catccacaa ggctgatccg

20

<210> 93  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 93  
tctgccatcc cacaaggctg

20

<210> 94  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 94  
catattttct tctccatggc

20

<210> 95  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 95  
aggttcgagt cttccgagat

20

<210> 96  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 96  
gaatgtaccc ctgatatggc

20

SEQUENCE LISTING

&lt;210&gt; 97

<211> 2372

<212> DNA

<213> Mus musculus

 $\langle 220 \rangle$ 

&lt;221&gt; CDS

 $\langle 222 \rangle \quad (190) \dots (1101)$ 

&lt;400&gt; 97

gcggcgqget ccgtagcgcg cgttcccget cgcceccage ttcccgggc tgcaggcaca 60

gccccgtccg gccgcgccgc tctctccct ccgcctccgg gacttttget ttcagttttc 120

ccaaagctgc cctcgacct tgcggaggac ggacgctgcc gtgggctcct ggcgcgcgcc 180

gtgggaacg atg acc gat gat cag gac tgt gct gcg gag ctg gaa aag gtg 231  
Met Thr Asp Asp Gln Asp Cys Ala Ala Glu Leu Glu Lys Val  
1 5 10

gat tct tcc agc gaa gac gga gtt gac gcc aag cca gac cgc tcc tct 279  
Asp Ser Ser Ser Glu Asp Gly Val Asp Ala Lys Pro Asp Arg Ser Ser  
15 20 25 30

atc atc tcc tct att ctc ttg aag aag aag aga aat gcc tct gcg ggc 327  
Ile Ile Ser Ser Ile Leu Leu Lys Lys Lys Arg Asn Ala Ser Ala Gly  
35 40 45

ccc gtc agg acc ggc cgg gac cga gtg ccc act tat ctg tac cgc atg 375  
Pro Val Arg Thr Gly Arg Asp Arg Val Pro Thr Tyr Leu Tyr Arg Met  
50 55 60

gat ttc cag aag atg ggt aaa tgc atc atc ata aac aac aag aac ttc 423  
Asp Phe Gln Lys Met Gly Lys Cys Ile Ile Ile Asn Asn Lys Asn Phe  
65 70 75

gac aaa gcg aca ggt atg gac gtc cgg aat ggg acg gac aaa gat gca 471  
Asp Lys Ala Thr Gly Met Asp Val Arg Asn Gly Thr Asp Lys Asp Ala  
80 85 90

ggg gcc ctc ttc aag tgc ttc caa aac ctg ggt ttt gaa gta acc gtc 519  
Gly Ala Leu Phe Lys Cys Phe Gln Asn Leu Gly Phe Glu Val Thr Val  
95 100 105 110

cac aat gac tgc tct tgt gca aag atg caa gat ctg ctt aga aaa gcc 567  
 His Asn Asp Cys Ser Cys Ala Lys Met Gln Asp Leu Leu Arg Lys Ala  
 115 120 125

tct gag gag gac cac agc aac tcg gcc tgc ttc gcc tgc gtc ctg ctg 615  
 Ser Glu Glu Asp His Ser Asn Ser Ala Cys Phe Ala Cys Val Leu Leu  
 130 135 140

agc cac ggg gaa gag gac ctg att tac ggg aaa gat ggc gtg aca ccc 663  
 Ser His Gly Glu Glu Asp Leu Ile Tyr Gly Lys Asp Gly Val Thr Pro  
 145 150 155

ata aag gat ctg aca gct cat ttt agg gga gac cga tgc aaa acc ctg 711  
 Ile Lys Asp Leu Thr Ala His Phe Arg Gly Asp Arg Cys Lys Thr Leu  
 160 165 170

tta gag aaa ccc aaa ctc ttc ttc att cag gca tgc cga ggg acg gag 759  
 Leu Glu Lys Pro Lys Leu Phe Phe Ile Gln Ala Cys Arg Gly Thr Glu  
 175 180 185 190

ctc gat gat gga atc cag gct gac tcg ggg ccc atc aac gac att gac 807  
 Leu Asp Asp Gly Ile Gln Ala Asp Ser Gly Pro Ile Asn Asp Ile Asp  
 195 200 205

gct aat ccc cgc aac aag atc ccg gtg gaa gcc gac ttc ctc ttt gct 855  
 Ala Asn Pro Arg Asn Lys Ile Pro Val Glu Ala Asp Phe Leu Phe Ala  
 210 215 220

tac tcc acg gtt cca ggt tat tac tca tgg agg aac cca ggg aaa ggc 903  
 Tyr Ser Thr Val Pro Gly Tyr Tyr Ser Trp Arg Asn Pro Gly Lys Gly  
 225 230 235

tcc tgg ttt gtg cag gcc ctc tgc tcc atc ctg aat gag cat ggc aag 951  
 Ser Trp Phe Val Gln Ala Leu Cys Ser Ile Leu Asn Glu His Gly Lys  
 240 245 250

gac ctc gag atc atg cag atc ctg acc agg gtg aac gac agg gtg gcc 999  
 Asp Leu Glu Ile Met Gln Ile Leu Thr Arg Val Asn Asp Arg Val Ala  
 255 260 265 270

agg cac ttc gag tcc cag tct gat gat cca cgc ttc aac gag aag aag 1047  
 Arg His Phe Glu Ser Gln Ser Asp Asp Pro Arg Phe Asn Glu Lys Lys  
 275 280 285

cag atc ccg tgt atg gtg tcc atg ctc acc aaa gag ctg tac ttc agc 1095  
 Gln Ile Pro Cys Met Val Ser Met Leu Thr Lys Glu Leu Tyr Phe Ser  
 290 295 300

cgt tga ccacccttca gctgagaacc tgccgccgtt cgttgatgaa tccagttttt 1151  
 Arg

attttatttt tgttccgatg ctctcaaaat atccagaaat gttgagggga ttttaatttca 1211

ggaaagtcta gatttttttt tttttgttta ataactttgt tcatctgatg acttcatgct 1271

cttctctata ggttgatttc ctgtttctgt ttcttttttt ctttgcgtc togctgagt 1331

catgctgtga gcatgacctc tggagaagac attggcaatg acgtctcagt tgaacttggc 1391

aaagagaatc ccagctcttg atgaaagaat acagctgcga cacctgttgg cctccattgg 1451

caaagggtgg tgctgagtgg ttgtttctcag tggttaggg cagattttta agccgacctt 1511

cccagggtgg tgagagaaga cgacagttaa tattccagta tatagaacct aatccagaaa 1571

ataagccatc ctaggaatat cgggtgcagaa ggggtcaatac agggaaattca cttactcggg 1631

taattgagag acagtcaatc ccgttacctc acatgcctct gtggggctgt aagaggactt 1691

cggttccctt gtcacagta agaactgtgg cagggcagcc ttctagagac tagaacctatg 1751

gaggactatg gtgattggag ccaggctcag gacaagccac tgccctgagat ggaaaccaag 1811

ccagaactga gtcagacacc aaaggcctcc ccaggcctgt gcccatatcc accagcgcct 1871

tataattcca aggtgcctga aagccaacta gtgagggcct agggaccatg ctaccctgtg 1931

ctacagagca gaggaagggga tagcaaagca ggggtagtgg tgggggtaaa cttggctcaa 1991

aatgtgaatt aaataaccat gtcctctggc gtgctacaat gtattcattc actattcact 2051

gatttgcag atcatccatc cacacaggtg ctgaagagta acccatttca ctttgtatca 2111

aagataatgt ttttgtactt caaatacatc tggaattctt tcaaatatcc caagattttt 2171

ttttctgaat aatcttttgt tacctctggt tctacaaatg acaatttagg cgaagatcct 2231



agcagtttct ttccaagcta tcattttctt tgggaattcgt tatacattcc tgtttttccc 2291  
acctaataaa tggttgacag atgttcctat ttattgatta aaatgtgttt cagaaaaaaaa 2351  
aaaaaaaaa aaaaaaaaaa a 2372

&lt;210&gt; 98

&lt;211&gt; 416

&lt;212&gt; DNA

&lt;213&gt; Mus musculus

&lt;400&gt; 98

aaaaaaaaagc tgccctcgac ccttgccgag gacggacgct gccgtgggct cctggccgcc 60  
gccgtgggaa cgatgaccga tgatcaggac tgtgctgcgg agctggaaaa ggtggattct 120  
tcacgcgaag acggagttga cgccaagcca gaccgctcct ctatcatctc ctctattctc 180  
ttgaagaaga agagaaatgc ctctgggggc cccgtcagga ccggccgggtg cgccgggtgcc 240  
cactggtggg gaccgcaggg aggtggcaag tgaggagagag gtgcatgggt ggagagaaga 300  
ggggcgctcg caaagcgagg tttgtggagg tccggaatgg gacggagaaa gatgcagggg 360  
cgcgcttcaa gtgctgccaa aacctgggtt ttgaagtaac cgtccacaat gactgc 416

&lt;210&gt; 99

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 99

gtgcctgcag cccgggaaag 20

&lt;210&gt; 100

&lt;211&gt; 20

&lt;212&gt; DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 100

tgaaagcaaa agtccggag

20

<210> 101

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 101

atcatcggtc atcgttccca

20

<210> 102

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 102

cgctttgtcg aagttcttgt

20

<210> 103

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 103

acctgtcgct ttgtcgaagt

20

<210> 104

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 104

gtccatacct gtcgctttgt

20

<210> 105

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 105

tttgtccgtc ccattccgga

20

<210> 106

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 106

ggaagcactt gaagagggcc

20

<210> 107

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 107

ctcagcagga cgcaggcgaa

20

<210> 108

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 108

cgtggctcag caggacgcag

20

<210> 109

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 109

atgagctgtc agatccttta

20

<210> 110

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 110

ccctaaaatg agctgtcaga

20

<210> 111

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 111

cgggtctccc taaaatgagc

20

<210> 112

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 112

tttgcacggt tctcccttaa

20

<210> 113

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 113

cagggttttg catcggtctc

20

<210> 114

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 114

ggccccgagt cagcctggat

20

<210> 115

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 115

accgtggagt aagcaaagag

20

<210> 116

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 116

ctggaaccgt ggagtaagca

20

<210> 117

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 117

ggcctgcaca aaccaggagc

20

<210> 118

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 118

agcagagggc ctgcacaaac

20

<210> 119

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 119

caggatggag cagagggcct

20

<210> 120

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 120

tcattcagga tggagcagag

20

<210> 121

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 121

gccatgctca ttcaggatgg

20

<210> 122

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 122

aggtccttgc catgctcatt

20

<210> 123

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 123

ggtcaggatc tgcgatgctc

20

<210> 124

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 124

ttcacccctgg tcaggatctg

20

<210> 125

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 125

tgctggttcac cctgggtcagg

20

<210> 126

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 126

ctggccaccc tgcgttcac

20

<210> 127

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 127

tcagactggg actcgaagtg

20

<210> 128

<211> 20

<212> DNA



<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 128

gtgagcatgg acaccataca

20

<210> 129

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 129

ctctttggtg agcatggaca

20

<210> 130

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 130

aaggggtggtc aacggctgaa

20

<210> 131

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 131

tctggatatt ttgagagcat

20

<210> 132

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 132

aacatttctg gatattttga

20

<210> 133

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 133

cttagaggaa gagcatgaag

20

<210> 134

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 134

ggtcatgctc acagcatgca

20

<210> 135

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 135

agcagccacc ttgccaatg

20

<210> 136

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 136

tgagaacaac cactcagcag

20

<210> 137

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 137

aaaatctgcc ctaagccact

20

<210> 138

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 138

ttctatatac tggaatatta

20

<210> 139

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 139

attccctgta ttgacccttc

20

<210> 140

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 140

taagtgaatt ccctgtattg

20

<210> 141

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 141

tctcaattac ccgagtaagt

20

<210> 142

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 142

tctagaaggc tgccttgcca

20

<210> 143

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 143

ttctagttctc tagaggctg

20

<210> 144

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 144

tgactcagtt ctgcttggt

20

<210> 145

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 145

gcgctggtgg atatgggcac

20

<210> 146

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 146

ttggctttca ggcacettgg

20

<210> 147

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 147

ctcactagtt ggctttcagg

20

<210> 148

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 148

ttcctaggcc ctctagtt

20

<210> 149

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 149

tgtagcacag ggtagcatgg

20

<210> 150

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 150

acgccagagg acatggttat

20

<210> 151

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 151

cagtgaatag tgaatgaata

20

<210> 152

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 152

actcttcagc acctgtgtgg

20

<210> 153

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 153

gaaatggggtt actcttcagc

20

<210> 154

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 154

tcctgtatac aaagtgaat

20

<210> 155

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 155

cagaggtaac caaagattat

20

<210> 156

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 156

agatcttcgc ctaaattgtc

20

<210> 157

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 157

aaataggaac atctgtcaac

20

<210> 158

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 158

gtatgaaaaat tcctgtgatt

20

<210> 159

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 159

tttccacctg ccaattctac

20

<210> 160

<211> 20

<212> DNA



<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 160

gtgaaagtgt gctactgttg

20

<210> 161

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 161

tgaaaactgc agatttgatg

20

<210> 162

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 162

atagccatgt aatataatgt

20

<210> 163

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 163

ttcaatcact gtagtctaag

20

<210> 164

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 164

ctgtacgaag aacttcaatc

20

<210> 165

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 165

ttgatggctg tacgaagaac

20

<210> 166

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 166

aaggaaacct tttcatgcct

20

<210> 167

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 167

attcaaagca gcagagaggg

20

<210> 168

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 168

acttgggctg tgaacattca

20

<210> 169

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 169

aacacttggg ctgtgaacat

20

<210> 170

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 170

ggactctgag aacacttggg

20

<210> 171

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 171

gggcagcaac tcagtttttg

20

<210> 172

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 172

ctgcgggtccc caccagtggg

20

<210> 173

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 173

gacctccaca aacctcgctt

20

<210> 174

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 174

cattccggac ctccacaaac

2